

Impact of climate change on waterborne diseases

Enzo Funari^(a), Maura Manganelli^(a) and Luciana Sinisi^(b)

^(a)Dipartimento di Ambiente e Connessa Prevenzione Primaria, Istituto Superiore di Sanità, Rome Italy

^(b)Istituto Superiore per la Protezione e la Ricerca Ambientale, Rome, Italy

Abstract. Change in climate and water cycle will challenge water availability but it will also increase the exposure to unsafe water. Floods, droughts, heavy storms, changes in rain pattern, increase of temperature and sea level, they all show an increasing trend worldwide and will affect biological, physical and chemical components of water through different paths thus enhancing the risk of waterborne diseases. This paper is intended, through reviewing the available literature, to highlight environmental changes and critical situations caused by floods, drought and warmer temperature that will lead to an increase of exposure to water related pathogens, chemical hazards and cyanotoxins. The final aim is provide knowledge-based elements for more focused adaptation measures.

Key words: climate change, waterborne diseases, microbial pathogens, chemical contaminants, toxic cyanobacteria.

Riassunto (*Effetto dei cambiamenti climatici sulle malattie trasmesse dall'acqua*). I cambiamenti climatici e del ciclo idrologico metteranno a rischio la disponibilità d'acqua e aumenteranno l'esposizione ad acqua contaminata. Le alluvioni, le siccità, le grandi tempeste, le variazioni nella frequenza ed intensità delle piogge, il riscaldamento e l'aumento del livello del mare crescono in ogni parte del mondo e influenzeranno le caratteristiche biologiche e chimico-fisiche dell'acqua attraverso diversi meccanismi, con il conseguente aumento del rischio di malattie trasmesse dall'acqua. L'analisi della letteratura disponibile, presentata in questo articolo, evidenzia i cambiamenti ambientali e le situazioni critiche causate da alluvioni, siccità e crescente riscaldamento che causeranno un aumento di esposizione a patogeni, inquinanti chimici e cianotossine, legati all'acqua. Lo scopo è di fornire gli elementi scientifici di base per misure di adattamento mirate.

Parole chiave: cambiamenti climatici, malattie trasmesse dall'acqua, microrganismi patogeni, contaminanti chimici, cianobatteri tossici.

INTRODUCTION

Climate variability and change may greatly influence human health [1], directly, as for instance drowning or trauma in extreme weather events, or indirectly, by altering the characteristics of the natural environments and habitats hence increasing the exposure of human populations to risk factors.

Extreme weather events are by far among the most destructive disasters known, whether their toll is measured in lives, damages of built environment, destruction of critical infrastructure, loss of properties and economic activities, irreversible contaminations, forced population displacement, short and long term diseases. According to EM-DAT [2] disastrous weather events database (criteria to be accounted for EM-DAT data base: a) at least 10 people killed; b) more than 100 hundred people affected; c) call for international assistance; d) declaration of state of emergency), the number of affected people in the UN-ECE Euro Region in the last two decades has increased of about 400% compared to previous decadal period. Up to the

first semester of 2008, as a consequence of adverse meteorological disasters, 38 million people required health assistance and basic survival needs such as safe shelter, medical assistance, a safe water supply and sanitation. EU accounted for 29 million of affected people with an economic loss of about 270 US\$ billion, the highest rate in the world of economic loss per capita. In particular, Italy and Germany suffered major damages from floods and storms due to high population and infrastructure density. The overall scenario in the western hemisphere is similar. In the US more than 700 billion US\$ in damages were estimated for the period 1980-2008, mostly due to hurricane, severe weather and non-tropical floods. Trends constantly increased worldwide and they're expected to do the same in the future since, as stated in the latest 2012 IPCC SREX Report [3], observed changes in climate extremes reflect the influence of anthropogenic climate changes in addition to natural climate variability, with changes in exposure and vulnerability influenced by both climatic and non climatic factors. Specifically about floods, it is

likely that the frequency of heavy precipitations or the proportion of total rainfall from heavy rainfalls will increase in the 21st century over many areas of the globe. This is particularly the case in the high latitudes and tropical regions, and in winter in the northern mid-latitudes. Heavy rainfalls associated with tropical cyclones are likely to increase with continued warming induced by enhanced greenhouse gas concentrations. There is high confidence that locations currently experiencing adverse impacts such as coastal erosion and inundation will continue to do so in the future due to increasing sea levels, all other contributing factors being equal [3].

In the temperate zone, climate change is predicted to decrease the number of rainy days, but to increase the average volume of each rainfall event [4]; as a consequence, drought-rewetting cycles may impact water quality as it enhances decomposition and flushing of organic matter into streams [5]. Flooding is the most common natural extreme weather event in the European Region [2]. Flooding may be caused by heavy rainfall, tidal surges and rapid snow melt. According to a database on floods in Europe, the most extreme flash floods are greater in magnitude in the Mediterranean countries than in the inner continental countries [6]. Coastal flooding related to increasing frequencies and intensities of storms and Sea Level Rise (SLR) is likely to threaten up to 1.6 million additional people annually in the EU alone [7]. Larger storm surges produced by extreme storms, combined with a rising sea level, could result in much higher rates of coastal erosion, which would in turn affect the levels of saline intrusion into coastal freshwater [8].

Other climatic factors affecting the hydrological regime are temperature and droughts, both of them projected to become worse.

About temperature, models predict with large confidence a substantial warming in temperature extremes by the end of the 21st century. Also, on a global scale, the frequency and magnitude of warm daily temperature extremes will increase, while cold extremes will decrease. The frequency of annual hottest days are projected to go from 1 in 20 years to 1 in 2 years, with an increase in the annual daily temperature of 2 to 5 °C by the end of 21st century, even if regional variations will often differ from the global changes [3].

These phenomena are going to affect many characteristics of water basins, as it has already happened. Atmospheric warming has been associated with an increase in surface water temperatures since the 1960s in Europe, North America and Asia (0.2-2 °C) [4]. In several lakes in Europe and Northern America, the water temperature increase has influenced the stratification period that has lengthened by 2-3 weeks [9]. In the European rivers Rhine and Meuse, an increase in the average summer water temperature of about 2 °C has been observed over the last three decades, with temperature peaks during the two severe droughts in 1976 and 2003, with

a pH increase (due to a decrease in CO₂ concentration) [10-11]. Computer models predict an increase of around 2 °C by 2070 in European lakes, although differences can be estimated, depending on lake characteristics and season [12-13]. The residence time in lakes with, at present, a short residence time, will probably increase by 92% in 2050 in summer and there will be a significant increase in temperature in the epilimnion and hypolimnion in shallow lakes [13]; however, on a long period, deepest lakes will be most sensitive to warming, due to their higher heat storage capacity and will experience highest winter temperature [14].

Also the ocean state has changed, in response to changed surface thermal conditions. The heat content of the World Ocean has increased since 1955, leading to sea level rise through thermal expansion, in addition to transfer of mass from glaciers, ice sheets and river runoff, due to changing hydrological regime. The waters at high latitudes (poleward of 50°N and 70°S) are fresher in the upper 500 m, while the subtropical latitudes in both hemispheres are characterized by increase in salinity. However, while there are many robust findings regarding the changed ocean state, key uncertainties still remain, making difficult projections for the future [15].

Droughts, which can be described as an unusual long period with little or no precipitation, are expected to increase in some areas. Model projections for the next 50-100 years indicate that climate change will reduce discharges to coastal waters in southern South America, western Australia, western and southern Africa, and in the Mediterranean Basin with consequences on salinities and nutrients and sediment delivered to the coast [16]. Salinity will tend to advance upstream, thereby altering the zonation of plant and animal species as well as the availability of freshwater for human use. Saltwater intrusion as a result of a combination of sea-level rise, decreases in river flows, unsustainable freshwater withdrawal and increased drought frequency are expected to modify physical and chemical components of estuarine-coastal environments with secondary impacts on phytoplankton community. In Central and Southern Europe and the Mediterranean region, in North and Central America, northeast Brazil and Southern Africa, droughts will likely intensify in the 21st century, due to reduced precipitation and/or increased evapotranspiration [3]. Projection for the rest of the world are still inconsistent, due to lack of data, or incapacity of the model to include all the different causes of dryness [3].

During dry periods, reduced groundwater recharge and increased water abstraction due to warmer temperatures may cause further water stress by reducing groundwater table levels. For these reasons in coastal areas, droughts can cause the intrusion of seawater into freshwater aquifers. In general freshwater contamination by seawater of only 5% is enough to rule out many important uses including drinking-water supply, irrigation of crops, parks and gardens, and

the well-being of groundwater- dependent ecosystems [17].

To identify the role of climate change on the spreading of waterborne diseases is made difficult by the simultaneous influence of other causes, like destruction of habitats, extensive travels and migrations of human populations, drug and pesticide resistance, urbanization and increased population density, and availability of health services [18].

The aim of this paper is to review the available literature to show the potential increase of the burden of WBDs resulting from climate changes and particularly from floods, increase of temperature and droughts, with regards to risk factors as microbial pathogens, chemicals, cyanotoxins.

MICROBIAL PATHOGENS

Waterborne pathogens of human and animal faecal origin include a high number of viruses, bacteria and parasitic protozoa. Also several naturally occurring microorganisms can be pathogenic to humans, as various species of *Vibrio* (gastroenteritis, diarrhoea and septicemia), *Pseudomonas aeruginosa* (skin and ear infections), *Legionella pneumophila* (Legionnaire's disease) and amoebae (encephalitis) [19].

Waterborne pathogens of concern for humans have the following characteristics:

- "are shed into the environment in high numbers, or are highly infectious to humans or animals at low doses" (*i.e.* cystis of protozoa);
- "can survive and remain infectious in the environment for long periods, or they are highly resistant to water treatment;
- some types of bacterial pathogens can multiply outside of a host under favorable environmental conditions" [20]. Pathogenic microorganisms of human

and animal faecal origin enter surface waters mainly by discharges of raw and treated wastewater and by runoff from the land. The reservoirs and routes of exposition are schematically depicted in *Figure 1*, after Hurst [21]. Some pathogens, like *V. cholerae*, hepatitis A virus and *Schistosoma* are restricted to tropical areas, while others like *Cryptosporidium* and *Campylobacter* are more diffused [22].

Heavy rainfall and floods

More frequent and intense heavy rainfall/floods will cause higher pathogen concentrations in natural waters which will generally be reflected in worse quality of drinking and bathing waters, crops and shellfish. Indeed, heavy rainfall/floods can cause over-flooding of sewage treatment plants, runoff of animal dejections and manure, re-mobilisation and redistribution of contaminated sediments [7, 23-27]. Since the diffusion of pathogens depends on hydrodynamic of surface water bodies it can be expected that floods and heavy rainfall, by speeding up water fluxes carrying pathogens, will counteract the natural pathogen inactivation in the environment by UV and temperature.

Enhanced environmental levels of pathogenic microorganisms may result in increased incidence of diseases and occurrence of new ones [28]. In general it is expected that zoonotic infections may expand due to an increased washing into water of wild animal and livestock faeces. A significant problem can be posed by an increased presence of different strains of enteric viruses in water bodies, as they are resistant to treatment in sewage treatment plants; bathing water receiving treated waters and seafood reared in receiving water bodies can represent important source of exposure to these pathogens [29]. Furthermore, viruses can be the unseen etiological pathogens responsible for human diseases even




Fig. 1 | Sources of waterborne pathogen. The non-enteric pathogens are naturally present in surface waters, which can, therefore, be a reservoir. (Modified from Hurst [21]).

when waters meet regulatory criteria for faecal contamination, based on conventional bacterial indicators, which are less resistant than viruses and decay much faster in natural environment [29]. Studies on cryptosporidiosis suggest that in the future more intense precipitation events may increase the saturation of soil profiles and mobilize infectious oocysts more often, significantly increasing the risk [30].

There are several examples of waterborne diseases outbreaks associated to excessive rainfall [31-43]. The largest reported waterborne disease outbreak in the United States, due to the presence of *Cryptosporidium* cysts in drinking water, occurred in Milwaukee in 1993 and was related to heavy rainfall and associated runoff and consequent contamination of Milwaukee lake, the source of the waterworks of the area. It resulted in the deaths of 54 people and more than 403 000 ill [37, 44]. Contamination of groundwater after flooding has been associated to additional disease outbreaks like *Acanthamoeba keratitis* in Iowa (USA) [22]. An outbreak of giardiasis in Montana (USA) was related to excess rainfall [39]. *Cryptosporidiosis* cases in England and Wales were positively associated with maximum river flow [45]. *Escherichia coli* O157:H7 and *Campylobacter jejuni* were responsible for a waterborne outbreak, causing 7 deaths, 65 hospitalization and more than 2300 cases of gastrointestinal illness in the Canadian town of Walkerton [46]. In this case, drinking water, supplied by shallow groundwater wells, turned out to be contaminated by a cattle manure from a local farm, following a period of intense spring rainfall, an event that is considered to happen once every 60 years [46].

In the US, Curriero *et al.* [38] reported a statistically significant association between excess rainfall and waterborne disease outbreaks over a long period of time and on a national scale. The study was based on 548 reported outbreaks in the United States from 1948 through 1994. The results indicated that 51% of waterborne disease outbreaks were preceded by rainfall events above the 90th percentile and that 68% were preceded by events above the 80th percentile. Outbreaks due to groundwater contamination were preceded by a 2-mo lag in rainfall accumulations whereas surface-water contamination showed the strongest association with excessive rainfall during the month of the outbreak [38].

In the European Union, in 2007, only 17 waterborne outbreaks were reported by eight countries, clearly indicating an under-reporting; they involved 10 912 cases, with 232 hospitalizations. The main microorganisms involved were *Campylobacter*, norovirus, *Giardia* and *Cryptosporidium* [47]. Only 24% of waterborne pathogen outbreaks in England, between 1970 and 2000, were found associated to heavy rainfall [48]. Yet, Nichols *et al.* [36] analyzing a small dataset [89] of waterborne outbreaks in England and Wales between 1910-1999, due to *Giardia*, *Cryptosporidium*, *E. coli*, *S. typhi*, *S. paratyphi*, *Campylobacter* and *Streptobacillus mon-*

iliformis, assessed a significant strong correlation between 40% of the cases and heavy rainfall in the week before the outbreak or low rainfall during the four weeks preceding the outbreaks.

Floods and hurricanes, destroying the water distribution system and mixing drinking and waste waters, can have a significant impact also on the diffusion of cholera, caused by the naturally occurring *V. cholerae*. The disease is one of the most severe forms of waterborne diarrheal disease, especially for developing countries, where outbreaks occur seasonally and are associated with poverty and use of poor sanitation and unsafe water [1]. The extreme climatic events increase the cases of diseases and fatalities by adding an oral-faecal contamination pathway difficult to manage.

Rainfalls are known to worsen the microbiological quality of bathing waters, indeed they are used directly as predictors of short term events of contamination in the European directive concerning the management of bathing water quality [49]. The same approach should be applied to shellfish-growing waters, as well. Indeed after heavy rainfalls sudden contamination of coastal waters are expected and results from environmental investigation, which require a too long time, do not represent the adequate tool to predict this contamination and prevent from dangerous human exposures. Heavy rainfall and sewage treatment plant failure were twice responsible for international gastroenteritis outbreaks due to consumption of oysters harvested from Tahu lagoon, in France [50, 51]. One of the outbreaks was characterized by the high diversity of human enteric viruses (up to six different strains) detected both in patient stool and shellfish samples [51]. Other important outbreaks associated to contaminated oyster/clam consumption caused by sewage overflow and discharge into the aquatic environment during heavy rainfall events were reported as those that affected 2000 people in Australia, in summer 1978 [52], and 1000 people in New York State in 1982 [53].

Temperature

There is a contrast between the well documented and forecasted increase in temperature and the paucity of data on the effects of this increment on microbial pathogens and infectious diseases [54].

Enteric pathogens in the water environment are generally neutralized by higher temperatures, however, their sensitivity shows different features. For instance, cysts of *Giardia* and enteroviruses are less rapidly inactivated compared with oocysts of *Cryptosporidium* [55]. *Cryptosporidium* oocysts are inactivated during the winter, as they are susceptible to freezing and thawing cycles; hence, as the number of frozen days decrease, oocysts may increasingly survive through the winter [30]. It is also known that a large variation in temperature susceptibility exists among viruses [56], suggesting a possible selection of more resistance strains, like the hepatitis A virus (HAV), which is fairly temperature insensitive. For some viruses a

correlation between increasing temperature and inactivation rate starts only after 10 days [55].

Increasing temperature could favor less temperature sensitive species, directly promoting the growth of some indigenous bacteria, including pathogenic species [57]. Vibriosis caused mainly by *Vibrio parahaemolyticus*, *V. vulnificus* and *V. alginolyticus* are among the 6 most common foodborne diseases monitored by the Foodborne Diseases Active Surveillance Network (FoodNet) in the USA, where *V. parahaemolyticus* in the past decade has become the main cause of gastroenteritis [58]. According to CDC, the incidence of *Vibrio* infections in the USA, which are also monitored by the Cholera and Other *Vibrio* Illness Surveillance (COVIS) system and are due to exposure to recreational water as well, was 76% higher in 2011 than in 1996-1998 [59]. A strong link between rising summer water temperatures, prolonged summer seasons, and noncholera *Vibrio* sp. infections has been shown [60], even if Johnson *et al.* [61] found that temperature explain only about 50% of the presence of *V. parahaemolyticus*, and also that the pathogenic subpopulations respond differently than the whole population to temperature.

The influence of increased temperatures on environmental bacteria is not expected to be homogeneous. The Baltic Sea for instance provides an environment in which only small changes of the present conditions (*e.g.*, temperature) result in increased *Vibrio* sp. populations [60].

Increasing temperatures would be expected to expand the range and increase the prevalence of *V. cholerae* and cholera both geographically and temporally, if environment and public health measures are not implemented [57]. Indeed, temperature shifts will alter the latitudinal distribution of planktonic species. Due to sea level rise inland areas will experience greater saltwater intrusion and increased levels of marine and estuarine bacteria, including *V. cholerae* [57]. It is then expected that an increase in temperature will threaten water quality with regard especially to cholera disease in Asia and South America [22]. However, the question may be more complex. A non linear population model to explain and predict the dynamics of *V. cholerae* has been developed by Koelle *et al.* [62]. The authors, considering both extrinsic (climatic) and intrinsic (acquired temporary immunity) drivers of epidemics, could explain the interannual cycles of cholera outbreaks from Matlab, Bangladesh. They found a strong correlation with climatic variables (monsoon) over 7 years and with local water temperature, degree of floods and droughts, at a shorter scale. Warmer pond and rivers increase the incidence of cholera through the faster growth rate of the pathogen in aquatic environments, but to have a good correspondance between the model and the incidence of cholera epidemics it was necessary to include in the model data on other parameters included the complex dynamics of acquired temporary immunity which could explain the interval between two epidemics [62].

There are also some authors who argued that one of the main effects of climate change is the reduction of biodiversity, that is some species are disappearing at a high rate, and that pathogens are subjected to the same ecological constraints [63]. Therefore, if there are areas where health conditions are worsening due to the spreading of pathogenic organisms, there are other areas that will be affected in the opposite way, thus reducing the areal distribution of the same pathogen. The final balance could be no overall increase in the global diffusion of a pathogenic species. The importance of such a debate is to highlight, among others, some important aspects to be considered: 1) better data sets and modeling approaches are required to make robust predictions of the impacts of climate change on disease dynamics and 2) expansion or reduction of specific pathogens geographical ranges will depend not only on extrinsic factors (including climate change), but also on intrinsic factors (such as immunity, phenotypic plasticity, and evolution) [64].

Droughts

The impacts of drought on human health due to shortage of water are dramatic and include deaths, malnutrition, increase in infectious diseases [65]. These effects are associated with worsening hygienic conditions, higher probability of microbial contamination of drinking water due to infiltration of organic material along the distribution system when pressure drops, higher re-use of wastewaters in agriculture, with consequent contamination of fresh vegetables, unsafe use of untreated water. Additionally, water shortages may increase the likelihood of multiple uses of a water body (*e.g.*, for cleaning, bathing, and drinking) with a consequent increase of the risk of microbial contamination and human exposure to pathogens [57]. Periods of droughts followed by short intense rainfall can cause peaks of surface water contamination. River-bed sediments represent an important microorganisms reservoir in dry areas, like the Mediterranean, where long dry periods are interrupted by flash-floods, transporting most of them downstream, up to coastal waters [66]. In areas affected by increasing droughts, treated effluents from sewage plants might become a quantitatively important source of water influx into river and heavy rainfall events can pose a risk to human health because of the huge increase of pathogenic microorganisms concentrations in the river downstream sewage treatment plants. In a study on *Campylobacter*, it has been shown that the resulting combined sewer overflow, enriched in pathogen, is discharged into the receiving rivers at a 150 fold higher concentration than usual [67].

Drought can lower the water table, resulting in changes in underground water flows of surface water into groundwater. In UK, an outbreak of *Cryptosporidium*, due to contamination of the borehole used as a source of drinking water was recorded when unusually very strong rainfall followed a long dry period [68]. No conclusive explanation is

given in the report, but the authors suggested the possible contamination of the aquifer by the intrusion of a contaminated river water flowing nearby, through interstices in the chalk [68]. Other examples of outbreaks linked to contamination of groundwater sources have been documented. In Brushy Creek, Texas, an outbreak of cryptosporidiosis was reported in 1998 followed extended drought conditions. The primary drinking water supply of Brushy Creek was chlorinated groundwater [69]. This source was contaminated by sewage, through fractures in the bedrock due to the long period of drought and extreme heat in which heavy water demand and no rainfall was present to recharge the aquifer [69]. Outbreaks of cryptosporidiosis have been associated to contamination of drinking water supplied by surface water, due to intense rainfall after very long and unusual drought periods, in Japan and Oregon [70-71]. The largest reported outbreak of *E. coli* O157:H7 occurred at a fairground in the state of New York in September 1999 and was linked to contaminated well water. This outbreak resulted from unusually heavy rainfall, which was preceded by a drought [43].

In general, monitoring programs are not elaborated in such a way to capture these periods of very high peaks of pathogenic concentrations which in turn represent a particularly high risk of infectious diseases transmission.

CHEMICALS

Climate change may influence the concentrations of chemical constituents and contaminants in natural waters through diverse ways. As an example, coastal erosion, likely to be exacerbated under climate change, has led to the exposure of landfill sites in Europe, with a clear potential for contamination of coastal waters [27].

It is difficult to estimate the risk associated to increased chemical concentrations in natural waters. Nevertheless, it seems possible to identify the following scenario of human exposure and the respective reasons of concern.

Heavy rainfall and floods

Climate change will influence the concentration of chemical constituents in natural waters. Indeed, it has been reported that a storm flow, by increasing the concentration of dissolved organic matter to which some metals can strongly get complexed, could lead to a transport of dissolved lead, titan and vanadium in peat land systems [72]. Furthermore a seasonal change in dissolved metal concentrations has also been observed for various trace elements (Fe, Mn, Al, La, U, Th, Cd and As). An increase of organic carbon content and a decline in redox conditions seem to be related with a trace elements release. A positive correlation is also found between storm events and trace element concentrations in streams [73]. In fact, organic and inorganic colloids could play an important role in trace elements mo-

bilization in soil sand water [74]. The possible implications for human health associated with these higher mobilization of chemical constituents from minerals and higher concentrations in natural waters are difficult to demonstrate. Natural waters contain many minerals, often in very small concentrations, that arise from the contact with the rocks and soils that water goes through. Most of these are of no concern but some are known they may impact human health for instance via drinking water, as arsenic and fluoride. Both arsenic and fluoride are significant contributors to morbidity in regions where concentrations in water are high. Hence, higher concentration of these elements as well as others of human concern may represent higher risky conditions in the affected areas.

More frequent intense rainfall is likely to exacerbate the flushing to water bodies of agricultural pollutants, including pesticides and veterinary medicines [28, 75]. Flooding may lead to contamination of water other than with chemicals already in the environment like pesticides, with dangerous chemicals, heavy metals, or other hazardous substances, from storage. Overall, flood events and strong rainfalls can transport pollutants from a contaminated area to a non-contaminated one [24, 76], from soil and sediments to water bodies.

Drinking water treatment is vital in protecting public health against microbial illness which remains a major cause of morbidity and mortality in many parts of the world, yet care has to be taken that treatment does not introduce higher levels of unwanted disinfection by-products (DBPs) than necessary. Indeed, epidemiological studies reported weak associations between chlorination and cancers of the colon, rectum and bladder and positive associations between concentrations of DBPs and a number of adverse reproductive effects, particularly stillbirth and low weight for gestational age [77]. In the past years a significant decline in DBPs concentrations has been achieved especially thanks to the introduction of improved treatment to remove the natural organic matter and better filtration that means less chlorine to be added. Floods and heavy rainfalls may increase concentrations of organic precursors of DBPs in surface waters [14]. On the other hand they cause higher nutrient concentrations which can promote algal and cyanobacterial blooms. Chen *et al.* [78] showed that DBPs precursors originating from high cyanobacterial densities can account for significant percentages of the total DBPs formation potential.

Increased floods and rainfalls will alter the transport, transfer, deposition and fate of chemical contaminants in coastal waters. Bioavailability of specific contaminants (*e.g.* metals) is greatly affected by salinity [79, 80]. Numerous studies have shown an increasing metal uptake by diverse aquatic organisms at reduced salinities [81-83]. Other studies have shown that solubility of many PAHs depends on salinity in the ambient water [84]. Thus, floods and

rainfalls will regulate the extent of exposure to toxic substances of seafood, hence of its consumers.

Unfortunately there is still little published evidence demonstrating a causal effect of chemical contamination on the pattern of morbidity following flooding events. But it is expected that floodings and heavy rainfalls, having implications for residue levels in food crops, food animals [85] and water bodies [86], will increase human exposure to chemical contaminants.

Temperature

It is well known that temperature influences physical chemical equilibria and biological reactions. Chemical reactions can be doubled for a temperature increase of 10 °C. As a consequence, higher water temperature will determine an increase of dissolution, solubilization, complexation, degradation, evaporation. Hence higher temperatures will lead to higher concentration of dissolved substances in water but will also favor an increase of volatilization of chemicals.

Considering the specific issue of pesticides, higher temperature will exert contrasting impacts, from one side higher degradation and volatilization [75] – which often will cause a more intensive use of them to fight pests risk – but from the other side, more solubilization in water. It is likely that changes in land use, induced by higher temperature, might play a more significant effect on pesticides in the environment than transformation and repartition processes [75]. For example, an increased prevalence of pests, weeds and diseases may lead to wider and more frequent application of both pesticides and veterinary medicines [28, 75].

Higher temperature in the Oceans may increase human exposure to mercury, especially in geographical areas where the population diet is based on seafood. Mercury is a global pollutant and is a reason of concern for public health when it is elevated above natural background levels, mainly through anthropogenic causes [87]. Mercury seems methylated by biotic processes [88]. Monomethyl mercury bioaccumulates and biomagnifies at all trophic levels in the food chain and can have severe neurological effects. Mercury methylation rates are temperature dependent [89]. Usually 80-99% of mercury found in fish muscle tissue is methyl mercury, regardless of its concentration in the environment [89]. It is expected that ocean temperature changes will increase the number of people exposed to above the tolerable weekly intakes defined by WHO for mercury [90].

Increased temperatures of natural surface waters used as drinking water supplies promote the DBPs formation rate [14]. Rodriguez and Serodes [91] showed that trihalomethanes (THM) concentrations vary from 1.5 to 2 times, depending on the utility, between drinking water plant and tap. In the same way, others authors reported that increasing temperature (10-33 °C) generally increased the formation of bromoorganic DBPs [92].

Droughts

Hotter, drier summers and increasingly severe and frequent droughts will deplete river flows, reducing contaminant dilution capacity and leading to elevated concentrations of hazardous substances [27]. Dry periods may entail brief spikes of compounds from sewage effluents, which can provoke transient perturbation of river ecosystems [93], with possible implications for human health. Increased contaminant concentrations are expected also in groundwater aquifers, especially in the unconfined ones.

Connected to water shortage is the particular case of temporary rivers. These are characterized by periodic dry phase and are a significant percentage of total river length in the world (between 40 and 70%). They are dominant in the semiarid Mediterranean area [94], where they represent an important source of water. These ecosystems are particularly exposed to the alteration of hydrological cycle due to longer drought periods associated with intense runoff and flushing [16], which can bring particularly high concentrations of pathogenic microorganisms and chemicals. The rising interest in their ecology during the last decade has highlighted their importance as links between water stored in soils, aquifers, snowpack, glaciers, vegetation and the atmosphere [95]. Recently, it has also been shown that they are important spots of nutrient and carbon recycling during the dry period [96, 97]. These processes are strictly related to the biochemistry of hazardous substances, entered the river along the hydrographic basin. Increasing frequency and duration of alternated dry-heavy rain periods, due to climate changes, by altering the timing of these processes is definitely going to affect the impact on the coastal environments of the catchments. Whether in a positive or negative way, there are not enough data yet [98].

CYANOBACTERIA

Health risks due to the toxins produced by cyanobacteria, photosynthetic prokaryotes diffuse in all the habitats, especially the aquatic ones, have been extensively addressed in this same issue of *Annali dell'Istituto Superiore di Sanità* by Manganelli *et al.*, p. 415 [99]. Briefly, cyanotoxins have different toxicological profiles, and target different organs. Several environmental factors affect their production, but mechanisms are still unknown. Humans can be exposed to cyanotoxins via ingestion (drinking and bathing water, through aquatic food chain), via aerosol and via parenteral, if surface contaminated water is used for haemodialysis [99]. Climate changes effects will be summed up with the effects of other environmental variables on cyanobacterial fitness and toxicity, with different outcomes, depending on the species/strain and the environment (*i.e.* lake, river, seawater). Available data are mostly obtained in laboratory studies or using mesocosm systems, even providing conflicting results: in many cases the apparent discrepancy could be attributed to the diverse response to climate change of differ-

ent cyanobacterial species, strongly depending on their physiological or ecological features, or, in the case of field studies, to the different response of the water body. It is therefore not possible to draw some general behaviours for the entire class of cyanobacteria and separated analyses should be carried out depending on the species and the environment. Bearing in mind these considerations, it is possible, at present, only summarize the available information in the following as shown in *Figure 2*.

Heavy rainfall and floods

Changes in freshwater runoff will have the greatest potential impacts on estuaries, causing changes in physical mixing characteristics [100]. Freshwater inflows into estuaries influence water residence time, nutrient delivery, vertical stratification, salinity, and control of phytoplankton growth rates [101]. Increased freshwaters inputs from increased runoff can dilute the estuarine environment by lowering salinity and promote extremely large blooms of toxic cyanobacteria by carrying *inocula* of salt-tolerant cyanobacteria from inland rivers. A combination of a particularly strong freshwater inflow after one day of unusual heavy rainfall with high temperature and the right nutrient concentration, favoring fast growth rate, has been the cause for a very large bloom of *Microcystis aeruginosa* in the Swan River Estuary, West Australia, in February 2000 [102].

Since increasing salinities can induce cell lyses and/or extracellular release of toxins [103], an indirect effect of heavy rainfall/floods, by conveying toxic cyanobacteria to brackish coastal waters, can be an increase in the extracellular quota of toxins. In 2005 in St. Lucie River Estuary, one of the largest brackish water systems on the East Coast of Florida, a periods of heavy rainfall caused the washout of *M.*

aeruginosa cells into the estuary, suddenly covered by a dense bloom: the abrupt change in salinity (32‰) caused an increase of 80% of toxin release in water [104]. Heavy rainfall and floods enhance the possibility of cyanobacteria expansion into coastal environments, thus increasing the exposure of edible organisms [99]. It is worth noting that heavy rain and floods can increase the nutrient availability of lakes, that in turn induce cyanobacterial proliferation. In Bangladesh, where a heavy rainfall drained phosphorus from the surrounding paddy fields into an aquaculture ponds, the increased nutrient concentration (9.5 mg L^{-1}) coincident with high temperature ($31 \text{ }^\circ\text{C}$) was the possible cause of a bloom of *Microcystis aeruginosa* and *Aphanizomenon flos-aquae* [105]. Soil runoff and discharge from wastewater treatment plants, rich in phosphorus, iron and carbon, might be responsible for the spreading of the mat-forming nitrogen-fixing *Lyngbya maiuscula* in coastal Queensland [106-107]. In Denmark the increase in phosphorus loading from land to lakes and coastal areas is expected to increase by 3 to 16% in the next 100 yr, due to higher winter rainfall; a shift in lakes community has already been observed towards dominance of cyanobacteria and dinoflagellates vs. diatoms and crysophytes [108].

Temperature

Increasing temperature in eutrophic condition can directly increase cyanobacterial proliferation and can be a main factor for their poleward movements and possibly their toxicity [109]. Indeed, some cyanobacteria like *Microcystis*, *Anabaena* and *Oscillatoria* have higher growth rate at higher temperature with respect to diatoms [110-112] and temperature can be one of the main factor in determining the success of one species vs other cyanobacteria or other phyto-




Fig. 2 | Schematic diagram depicting the interrelations between changing meteorological parameters and toxic cyanobacteria.

plankton, if nutrients concentration is not limiting [113-115].

It is expected that the warming trend will move forward the onset of spring bloom in high latitude and temperate areas [116] and will accelerate the spread and abundance of subtropical species to temperate regions all over the world. *Cylindrospermopsis raciborskii* is probably the best known example. It was originally found in the tropics, in central Africa and Australian lakes; at present, starting from the mid 90s, it has been reported in several temperate areas, in Europe and central America. Climate change is one of the factors considered in explaining its diffusion [117, 118]. Notwithstanding the areal expansion, the risk due to the exposure to *C. raciborskii* cannot be directly inferred and it is not straightforward that it will increase, since the toxins produced in different continents are different: the Australian strains produce cylindrospermopsin [119], whereas the Brazilian or European strains produce saxitoxins and a still unknown toxin, respectively [120, 121].

Another example of expected larger diffusion of tropical species is that of *Trichodesmium* spp. in seawater. Temperature dependence of O_2 flux and respiration seems to be the main reason for this non heterocystous N_2 fixing species to be dominant in subtropical and tropical ocean regions [122]. The reduced O_2 flux in warm seawater and the high respiration rate at high temperature allow *Trichodesmium* to outcompete other heterocystous species. It has been estimated that the poleward shift of the 20 °C isothermal as a consequence of a 3 °C increase by 2090, which poses a lower limit to the distribution of *Trichodesmium*, will expand its range of 11%, even if this will be counteracted by the reduction of 16% of its thermal niche at the tropics, where temperature is expected to exceed 30 °C, corresponding to suboptimal growth rates [123]. It is therefore reasonable to expect a poleward shift of this species, being all the other parameters unchanged.

Complex interactions between direct and indirect effects of temperature are more often the cause of shifts in community composition and changes in dominant species. Increasing thermal stability and stratification in deep, eutrophic lakes will favor the proliferation of buoyant cyanobacteria, like *Microcystis*, *Anabaena*, *Aphanizomenon*, and elevate the chance for the development of surface blooms [124]. Milder winter in temperate eutrophic lakes, may indirectly favor also Oscillatoriales bloom, by inducing a complex shift in phytoplankton community composition [125].

The observed significant increase in cyanobacterial biomass over the years in the Baltic Sea, composed mainly by *Aphanizomenon flos-aquae*, *Nodularia spumigena* and *Anabaena* sp., has been correlated to increased temperature, to variation in the hydrology, and to nutrients inputs [126]. Hence a general higher exposure of aquatic organisms, and possibly of their consumers, human beings included, can be foreseen.

Data available indicate that the prolonged thermal stratification, due to warming climate, would favor also species like *Planktothrix rubescens* [127], which are usually more adapted to colder water.

Data on increasing temperature effects on toxicity are controversial, and depend on the co-occurrence of other parameters, like light and nutrients [128-129]. Davis *et al.* [129] showed that in 4 lakes in Northeast US, higher temperature (and phosphorus concentration) favored an increase of microcystins MCs production by stimulating the growth rate of *Microcystis* spp. toxic cells [129]. During the last decade, the rise of minimal surface temperature of many reservoirs and water basin in South Africa seems to be the direct cause of the proliferation and bloom of toxic cyanobacteria, since there has been a correlated increase of animal mortalities in several areas [130].

In *Oscillatoria agardhii* isolated from Finnish lakes MCs production rate peaked at optimum growth temperature in one strain (25 °C) and was constant in the range 15-25 °C in a second one. However, for both strains, growth and toxin production decreased at 30 °C [131]. On the contrary, for *Microcystis aeruginosa* increasing temperature from 18 °C to 28 °C, yielded higher growth rates, but reduced toxicity of the cultured strains [132]. Also the production of *cylindrospermopsin* CYN seems to be inhibited by higher temperature in tropical *Cylindrospermopsis* [119] and in temperate *Aphanizomenon* [133].

Additionally, temperature can impact on toxicity by affecting the variants profile of toxin production, as it happened in an *Aphanizomenon* sp. cultured strain, which following an increase of T from 22 to 28 °C, switched towards the production of the more toxic variant among the 4 detected toxins [134].

In two strains of *Anabaena* from a lake in Southern Finland, temperature higher and lower than the optimum (25 °C) decreased the total amount of MCs production. In addition the profile of MCs variants changed: high light and high temperature favored the production of the less toxic -RR variant, while an increase in -LR variant was observed at lower temperature [135].

Droughts

The likely longer dryness periods, increasing in some regions at mid-latitudes and in the dry tropics, some of which already water-stressed areas, may influence cyanobacterial proliferation by increasing nutrient availability (higher concentrations due to surface water evaporation in summer) and reducing water bodies flow (thus increasing the areas of still waters in which cyanobacterial growth is easier). One of the largest bloom of *Anabaena circinalis* involved more than 1000 km of one of Australia's major river systems, in a very low flow conditions, concurrent with high nutrients concentration and high temperature [136].

Another important consequence of droughts is that rainfall following period of dryness, can transport cyanotoxins into groundwater, beneath parched and cracked soil, thus contaminating very precious sources of drinking water.

Changes in phytoplankton communities of coastal environment exposed to a reduction in freshwater discharge and to saltwater intrusion, can eliminate more sensitive species and favor more tolerant cyanobacteria and expose aquatic organisms to higher concentration of dissolved cyanotoxins [103].

In Nebraska lakes a lower water quality due to the presence of MCs, associated with the drought conditions and lower nitrogen:phosphorus ratios [137], was reported during 2004, concomitant with dogs, wildlife and livestock deaths, and more than 50 accounts of human skin rashes, lesions, or gastrointestinal illnesses.

CONCLUSIVE REMARKS

Reviewed data show that increase of water temperature, heavy rains, floods and droughts will increase the distribution and patterns of human exposures to pathogens, chemicals and cyanobacteria. Several studies have shown the association between heavy rainfalls/floods and outbreaks of waterborne diseases. The number and severity of these outbreaks are destined to increase. In particular, zoonotic infections will likely increase as a consequence of washing into water of wild animal and livestock faeces due to heavy rain falls and floods. Different strains of enteric viruses and protozoa will increase their concentrations in water bodies representing etiological pathogens not signaled by conventional faecal contamination indicators.

Several outbreaks have been reported due to groundwater contamination by pathogens as a consequence of drought, that has lowered the water table, resulting in changes in underground water flows of surface water into groundwater.

Increasing temperature could favor the growth of some pathogenic indigenous bacteria, like *Vibrio parahaemolyticus*, *V. vulnificus* and *V. alginolyticus*, which are already among the most common etiological agents responsible for diseases transmitted by seafood consumption in US. Increasing temperatures are expected to expand the range and increase the prevalence of *V. cholerae* and cholera, whose diffusion will also be favored by the destructive effects of floods and hurricane on water distribution systems.

Heavy rainfalls and floods will increase the release of chemical constituents from minerals and their concentrations in natural waters, including that of elements like As and F, that at high levels are dangerous for human health. Similarly they will increase concentrations of precursors of DBPs, leading to higher exposures through drinking water consumption. Higher temperature in the oceans may increase human exposure to methyl mercury,

especially in geographical areas where the population diet is based on seafood, increasing the number of people above the tolerable weekly intake defined by WHO.

Increased temperatures and floods will promote the formation rate of DBPs.

Heavy rainfalls and floods will expand the distribution of freshwater cyanobacteria into brackish and coastal waters, by physical transport of cyanobacterial blooms into estuaries and by altering the chemical and physical conditions of those areas, ultimately affecting the composition of phytoplankton. These effects will increase the risk of exposure of seafood to cyanotoxins, which is going to be one of the new scenarios of exposure.

Also temperature will affect the geographical distribution of tropical toxic species, that will move towards temperate latitudes, where water temperature is expected to increase. Many deep eutrophic lakes are likely to be dominated by buoyant cyanobacteria, by the protraction of water stratification, which will prolong the period of exposure.

Of course the above predictions do not consider the role of management measures that can be implemented to counteract the worsening scenario, especially to mitigate the effects of extreme events. For these factors, the main focus is identifying regions most at risk of flooding and preparing plans for responding and mitigating the main consequences, as stated in the European directive on management of floods [138].

Many Countries worldwide are strengthening their ability to cope with these events improving their early warning systems, vulnerability assessment and response plan to emergency, still we need a more coordinated approach among different operators (environmental, public and veterinary health, water managers, utilities sectors, land use managers) to counteract water unsafety.

WBDs health surveillance, post extreme events environmental and biota monitoring of affected areas, efficiency of waste water treatment are also crucial areas of action for adaptation measures as well as training of professionals involved at local level on new risk scenarios of WBDs.

If the increase of temperature and extreme events is a result of global warming due to human activities, losses, damages, increased burden of diseases could continue rising indefinitely. Long term measures to counteract the impact of climate change on human health rely on a sound strategy aimed at reducing green house gases emissions.

Conflict of interest statement

There are no potential conflicts of interest or any financial or personal relationships with other people or organizations that could inappropriately bias conduct and findings of this study.

Submitted on invitation.

Accepted on 24 September 2012.

References

- World Health Organization - World Meteorological Organization. *Atlas of health and climate*. Geneva: WHO, WMO; 2012. Available from: http://www.wmo.int/ebooks/WHO/Atlas_EN_web.pdf.
- EM-DAT. *EM-DAT international disaster database*. Brussels: Universite Catholique de Louvain Centre for Research on the Epidemiology of Disasters (CRED); 2010. Available from: www.emdat.be/database.
- Field CB, Barros V, Stocker TF, Qin D, Dokken DJ, Ebi KL, et al. (Ed.). IPCC 2012. *Managing the risks of extreme events and disasters to advance climate change adaptation. A special report of the intergovernmental panel on climate change*. Cambridge, UK, and New York, NY, USA: Cambridge University Press; 2012.
- Bates BC, Kundzewicz ZW, Wu S, Palutikof JP (Ed.). *Climate change and water. Technical paper of the intergovernmental panel on climate change*. Geneva: IPCC Secretariat; 2008.
- Evans C, Monteith D, Cooper D. Long-term increases in surface water dissolved organic carbon: observations, possible causes and environmental impacts. *Env Poll* 2005;137:55-71. <http://dx.doi.org/10.1016/j.envpol.2004.12.031>
- Gaume E, Bain V, Bernardara P, Newinger O, Barbuc M, Bateman A, et al. A compilation of data on European flash floods. *J Hydrol* 2009;367(1-2):70-8. <http://dx.doi.org/10.1016/j.jhydrol.2008.12.028>
- Menne B. Impacts of climate change and extreme events on waterborne disease. In: Sinisi L, Aertgeerts R (Ed.). *Guidance on water supply and sanitation in extreme weather events*. Copenhagen: UNECE/WHO; 2011.
- OzCoasts. *Saline intrusion*. Canberra ACT: OzCoast Australian Online Coastal Information; 2010. Available from: www.ozcoasts.org.au/indicators/saline_intrusion.jsp.
- Komatsu E, Fukushima T, Harasawa H. A modeling approach to forecast the effect of long-term climate change on lake water quality. *Ecol Modell* 2007;209(2-4):351-66. <http://dx.doi.org/10.1016/j.ecolmodel.2007.07.021>
- van Vliet MTH, Zwolsman JJG. Impact of summer droughts on the water quality of the Meuse river. *J Hydrol* 2008;353(1-2):1-17. <http://dx.doi.org/10.1016/j.jhydrol.2008.01.001>
- Zwolsman JJG, Bokhoven AJv. Impact of summer droughts on water quality of the Rhine River - a preview of climate change? *Anglais* 2007;56(4):45-55. <http://dx.doi.org/10.2166/wst.2007.535>
- Malmaeus JM, Blenckner T, Markensten H, Persson I. Lake phosphorus dynamics and climate warming: A mechanistic model approach. *Ecol Modell* 2006;190(1-2):1-14. <http://dx.doi.org/10.1016/j.ecolmodel.2005.03.017>
- George G, Hurley M, Hewitt D. The impact of climate change on the physical characteristics of the larger lakes in the English Lake District. *Freshwater Biol* 2007;52(9):1647-66. <http://dx.doi.org/10.1111/j.1365-2427.2007.01773.x>
- Delpla I, Jung AV, Baures E, Clement M, Thomas O. Impacts of climate change on surface water quality in relation to drinking water production. *Environ Int* 2009;35(8):1225-33. <http://dx.doi.org/10.1016/j.envint.2009.07.001>
- Bindoff NL, Willebrand J, Artale V, Cazenave A, Gregory J, Gulev S, et al. Observations: oceanic climate change and sea level. In: Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, et al. (Ed.). *Climate change 2007: The physical science basis contribution of working group I to the fourth assessment report of the intergovernmental panel on climate change*. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press; 2007.
- Milly PCD, Dunne KA, Vecchia AV. Global pattern of trends in streamflow and water availability in a changing climate. *Nature* 2005;438:347-50. <http://dx.doi.org/10.1038/nature04312>
- UNSW. *Potential impacts of sea-level rise and climate change in coastal aquifers*. Sydney, NSW: University of New South Wales; 2010. Available from: www.connectedwaters.unsw.edu.au/resources/articles/coastal_aquifers.html.
- Semenza JC, Menne B. Climate change and infectious diseases in Europe. *Lancet Infect Dis* 2009;9(6):365-75. [http://dx.doi.org/10.1016/S1473-3099\(09\)70104-5](http://dx.doi.org/10.1016/S1473-3099(09)70104-5)
- World Health Organization. *Guidelines for drinking-water quality: incorporating 1st and 2nd addenda*. Geneva: WHO; 2008.
- Rosen B. *Waterborne pathogens in agricultural watersheds*. Burlington, Vt.: USDA-NRCS Watershed Institute, University of Vermont; 2000.
- Hurst CJ. Overview of water microbiology as it relates to public health. In: Hurst CJ, Crawford RL, Garland JL, Lipson DA, Mills AL, Stetzenbach LD (Ed.). *Manual of environmental microbiology*: Third edition. Washington, DC: ASM Press; 2007. p. 219-21.
- Hunter P. Climate change and waterborne and vector-borne disease. *J Appl Microbiol* 2003;94:37-46. <http://dx.doi.org/10.1046/j.1365-2672.94.s1.5.x>
- Nie L, Lindholm O, Braskerud B. Urban flood management in a changing climate. *J Water* 2009;2:203-13.
- Hilscherova K, Dusek L, Kubik V, Cupr P, Hofman J, Klanova J, et al. Redistribution of organic pollutants in river sediments and alluvial soils related to major floods. *J Soil Sed* 2007;7(3):167-77. <http://dx.doi.org/10.1065/jss2007.04.222>
- Nagels JW, Davies-Colley RJ, Donnison AM, Muirhead RW. Faecal contamination over flood events in a pastoral agricultural stream in New Zealand. *Water Sci Technol* 2002;45(12):45-52.
- Muirhead RW, Davies-Colley RJ, Donnison AM, Nagels JW. Faecal bacteria yields in artificial flood events: quantifying in-stream stores. *Water Res* 2004;38(5):1215-24. <http://dx.doi.org/10.1016/j.watres.2003.12.010>
- European Environment Agency. *Hazardous substances in Europe's fresh and marine waters. An overview*. Luxembourg: EEA; 2011. (Technical report n. 8/2011).
- Boxall AB, Hardy A, Beulke S, Boucard T, Burgin L, Falloon PD, et al. Impacts of climate change on indirect human exposure to pathogens and chemicals from agriculture. *Environ Health Perspect* 2009;117(4):508-14. <http://dx.doi.org/10.1289/ehp.0800084>
- Maalouf H, Pommepuy M, Le Guyader F. Environmental conditions leading to shellfish contamination and related outbreaks. *Food Environ Virol* 2010;2(3):136-45. <http://dx.doi.org/10.1007/s12560-010-9043-4>
- King BJ, Monis PT. Critical processes affecting *Cryptosporidium* oocyst survival in the environment. *Parasitology* 2007;134:309-23. <http://dx.doi.org/10.1017/S0031182006001491>
- Howe AD, Forster S, Morton S, Marshall R, Osborn KS, Wright P, et al. Cryptosporidium oocysts in a water supply associated with a cryptosporidiosis outbreak. *Emerging Infect Dis* 2002;8:619-24.
- Kistemann T, Claßen T, Koch C, Dangendorf F, Fischeder R, Gebel J, et al. Microbial load of drinking water reservoir tributaries during extreme rainfall and runoff. *Appl Environ Microbiol* 2002;68(5):2188-97. <http://dx.doi.org/10.1128/aem.68.5.2188-2197.2002>

33. Auld H, MacIver D, Klaassen J. Heavy rainfall and waterborne disease outbreaks: the walkerton example. *J Toxicol Environ Health* 2004;67(20):1879-87. <http://dx.doi.org/10.1080/15287390490493475>
34. Thurston-Enriquez JA, Gilley JE, Eghball B. Microbial quality of runoff following land application of cattle manure and swine slurry. *J Water Health* 2005;3:157-71.
35. Mons C, Dumetre A, Gosselin S, Galliot C, Moulin L. Monitoring of *Cryptosporidium* and *Giardia* river contamination in Paris area. *Water Research* 2009;43:211-7. <http://dx.doi.org/10.1016/j.watres.2008.10.024>
36. Nichols G, Lane C, Asgari N, Verlander NQ, Charlett A. Rainfall and outbreaks of drinking water related disease in England and Wales. *J Water Health* 2009;7:1-8. <http://dx.doi.org/10.2166/wh.2009.143>
37. Mac Kenzie WR, Hoxie NJ, Proctor ME, Gradus MS, Blair KA, Peterson DE, et al. A massive outbreak in Milwaukee of cryptosporidium infection transmitted through the public water supply. *New Engl J Med* 1994;331(3):161-7. <http://dx.doi.org/10.1056/NEJM199407213310304>
38. Curriero FC, Patz JA, Rose JB, Lele S. The association between extreme precipitation and waterborne disease outbreaks in the United States, 1948-1994. *Am J Public Health* 2001;91(8):1194-9. <http://dx.doi.org/10.2105/AJPH.91.8.1194>
39. Weniger BG, Blaser MJ, Gedrose J, Lippy EC, Juranek DD. An outbreak of waterborne giardiasis associated with heavy water runoff due to warm weather and volcanic ashfall. *Am J Public Health* 1983;73(8):868-72. <http://dx.doi.org/10.2105/AJPH.73.8.868>
40. Lisle JT, Rose JB. Cryptosporidium contamination of water in the USA and UK: A minireview. *J Water SRT - Aqua* 1995;44:103-17. <http://dx.doi.org/10.1289/ehp.00108367>
41. Alterholt TB, LeChevalier MW, Norton WD, Rosen JS. Effect of rainfall on giardia and crypto. *J Am Water Works Assoc* 1998;90:66-80.
42. Rose JB, Daeschner S, Easterling DR, Curriero FC, Lele S, Patz JA. Climate and waterborne disease outbreaks. *J Am Water Works Assoc* 2000;9:77-87.
43. Patz JA, McGeehin MA, Bernard SM. The potential health impacts of climate variability and change for the United States: Executive summary of the report of the health sector of the U.S. National Assessment. *Environ Health Perspect* 2000;108:367-76.
44. Hoxie NJ, Davis JP, Vergeront JM, Nashold RD, Blair KA. Cryptosporidiosis-associated mortality following a massive waterborne outbreak in Milwaukee, Wisconsin. *Am J Public Health* 1997;87(12):2032-5. <http://dx.doi.org/10.2105/AJPH.87.12.2032>
45. Lake IR, Bentham G, Kovats RS, Nichols GL. Effects of weather and river flow on cryptosporidiosis. *J Water Health* 2005;3(4):469-74.
46. O'Connor DR. *Report of the Walkerton Inquiry. Part 1. The events of may 2000 and related issues*. Toronto: 2002.
47. European Food Safety Authority - European Centre for Disease Prevention and Control. The community summary report. Food-borne outbreaks in the European Union in 2007. Brussels, Belgium: EFSA-ECDC; 2009.
48. Said B, Wright F, Nichols GL, Reacher M, Rutter M. Outbreaks of infectious disease associated with private drinking water supplies in England and Wales 1970-2000. *Epidemiol Infect* 2003;130:469-79. <http://dx.doi.org/10.1017/S0950268803008495>
49. European Union. Directive 2006/7/EC concerning the management of bathing water quality and repealing Directive 76/160/EC. *Official Journal of the European Union* L 64 4/3/2006.
50. Doyle A, Barataud D, Gallay A, Thiolet J, Le Guyaguer S, Kohli E, et al. Norovirus foodborne outbreaks associated with the consumption of oysters from the Etang de Thau, France, December 2002. *Euro Surveill* 2004;9(3):451.
51. Le Guyader FS, Le Saux J-C, Ambert-Balay K, Krol J, Serais O, Parnaudeau S, et al. Aichi virus, norovirus, astrovirus, enterovirus, and rotavirus involved in clinical cases from a French oyster-related gastroenteritis outbreak. *J Clin Microbiol* 2008;46(12):4011-7. <http://dx.doi.org/10.1128/jcm.01044-08>
52. Murphy AM, Grohmann GS, Christopher PJ, Lopez WA, Davey GR, Millsom RH. An Australia-wide outbreak of gastroenteritis from oysters caused by norwalk virus. *Med J Aust* 1979;2(7):329-33.
53. Morse DL, Guzewish JJ, Hanrahan JP, Stricof R, Shayegani M, Deibel R, et al. Widespread outbreaks of clam- and oyster-associated gastroenteritis. Role of norwalk virus. *N Engl J Med* 1986;314(11):678-81. <http://dx.doi.org/10.1056/NEJM198603133141103>
54. Semenza JC, Höser C, Herbst S, Rechenburg A, Suk JE, Frechen T, et al. Knowledge mapping for climate change and food- and waterborne diseases. *Crit Rev Environ Sci Technol* 2011;42(4):378-411. <http://dx.doi.org/10.1080/10643389.2010.518520>
55. Schijven JF, de Roda Husman AM. Effect of climate changes on waterborne disease in The Netherlands. *Water Sci Technol* 2005;51(5):79-87.
56. Schijven JF, Hassanizadeh SM. Removal of viruses by soil passage: Overview of modeling, processes, and parameters. *Crit Rev Environ Sci Technol* 2000;30(1):49-127. <http://dx.doi.org/10.1080/10643380091184174>
57. Lipp EK, Huq A, Colwell RR. Effects of global climate on infectious disease: the cholera model. *Clin Microbiol Rev* 2002;15(4):757-70. <http://dx.doi.org/10.1128/cmr.15.4.757-770.2002>
58. DePaola A, Jones JL, Woods J, Burkhardt W, Calci KR, Krantz JA, et al. Bacterial and viral pathogens in live oysters: 2007 United States market survey. *Appl Environ Microbiol* 2010;76(9):2754-68. <http://dx.doi.org/10.1128/aem.02590-09>
59. Centers for Disease Control and Prevention. *A Trends in foodborne illness in the United States, 1996-2011*. Atlanta, GA: CDC; 2011. Available from: www.cdc.gov/foodnet/data/trends/trends-2011.html.
60. Semenza JC, Herbst S, Rechenburg A, Suk JE, Höser C, Schreiber C, et al. Climate change impact assessment of food- and waterborne diseases. *Crit Rev Environ Sci Technol* 2011;42(8):857-90. doi:10.1080/10643389.2010.534706.
61. Johnson CN, Flowers AR, Noriega III NF, Zimmerman AM, Bowers JC, DePaola A, et al. Relationships between environmental factors and pathogenic vibrios in the Northern Gulf of Mexico. *Appl Environ Microbiol* 2010;76(21):7076-84. <http://dx.doi.org/10.1128/AEM.00697-10>
62. Koelle K, Rodo X, Pascual M, Yunus M, Mostafa G. Refractory periods and climate forcing in cholera dynamics. *Nature* 2005;436(7051):696-700. <http://dx.doi.org/10.1038/nature03820>
63. Lafferty KD. The ecology of climate change and infectious diseases. *Ecology* 2009;90(4):888-900. <http://dx.doi.org/10.1890/08-0079.1>
64. Wilson K. Climate change and the spread of infectious ideas1. *Ecology* 2009;90(4):901-2. <http://dx.doi.org/10.1890/08-2027.1>
65. Menne B, Bertollini R. The health impacts of desertification and drought. *Down to Earth* 2000;14:4-6.
66. Chu Y, Salles C, Tournoud MG, Got P, Troussellier M, Rodier C, et al. Faecal bacterial loads during flood events

- in Northwestern Mediterranean coastal rivers. *Journal of Hydrology* 2011;405(3-4):501-11.
<http://dx.doi.org/10.1016/j.jhydrol.2011.05.047>
67. Rechenburg A, Kistemann T. Sewage effluent as a source of campylobacter sp. in a surface water catchment. *Int J Environ Health Res* 2009;19(4):239-49.
<http://dx.doi.org/10.1080/09603120802460376>
 68. Willocks L, Crampin A, Milne L, Seng C, Susman M, Gair R, et al. A large outbreak of cryptosporidiosis associated with a public water supply from a deep chalk borehole. *Communicable Disease and Public Health* 1998;1(4):239-43.
 69. Bergmire-Sweat D, Morgan J, Wilson K, VonAlt K, Marengo L, Bennett T, et al. (Ed.). *Cryptosporidiosis at Brushy Creek: describing the epidemiology and causes of a large outbreak in Texas, 1998*. International Symposium on Waterborne Pathogens. Milwaukee, WI: American Water Works Association; 1999.
 70. Leland D, McAnulty J, Keene W, Stevens G. A cryptosporidiosis outbreak in a filtered water supply. *J Am Water Works Assoc* 1993; 85(6):34-42.
 71. Yamamoto N, Urabe K, Takaoka M, Nakazawa K, Gotoh A, Haga M, et al. Outbreak of cryptosporidiosis after contamination of the public water supply in Saitama Prefecture, Japan in 1996. *Kansenshogaku Zasshi - J Japan Assoc Infect Dis* 2000;74(6):518-26.
 72. Rothwell J, Evans M, Daniels S, Allott T. Baseflow and stormflow metal concentrations in streams draining contaminated peat moorlands in the Peak District National Park (UK). *J Hydrol* 2007;341:90-104.
<http://dx.doi.org/10.1016/j.jhydrol.2007.05.004>
 73. Olivie-Lauquet G, Gruau G, Dia A, Riou C, Jaffrezic A, Henin O. Release of trace elements in wetlands: role of seasonal variability. *Water Res* 2001;35(4):943-52.
[http://dx.doi.org/10.1016/S0043-1354\(00\)00328-6](http://dx.doi.org/10.1016/S0043-1354(00)00328-6)
 74. Pédrot M, Dia A, Davranche M, Bouhnik-Le Coz M, Henin O, Gruau G. Insights into colloid mediated trace element release at the soil/water interface. *J Colloid Interface Sci* 2008; 325:187-97.
<http://dx.doi.org/10.1016/j.jcis.2008.05.019>
 75. Bloomfield J, Williams R, Goody D, Cape J, Guha P. Impacts of climate change on the fate and behaviour of pesticides in surface and groundwater-a UK perspective. *Sci Total Environ* 2006;369:163-77.
<http://dx.doi.org/10.1016/j.scitotenv.2006.05.019>
 76. Harmon S, Wyatt D. Evaluation of post Katrina flooded soils for contaminants and toxicity to the soil invertebrates *Eisenia fetida* and *Caenorhabditis elegans*. *Chemosphere* 2008;70(10):1857-64.
<http://dx.doi.org/10.1016/j.chemosphere.2007.08.007>
 77. Fawell J. Chemicals in the water environment. Where do the real and future threats lie? *Ann Ist Super Sanità* 2012;48(4):347-53.
 78. Chen C, Zhang X-J, Zhu L-x, Liu J, He W-j, Han H-d. Disinfection by-products and their precursors in a water treatment plant in North China: seasonal changes and fraction analysis. *Sci Total Environ* 2008;397:140-7.
<http://dx.doi.org/10.1016/j.scitotenv.2008.02.032>
 79. McLusky DD, Bryant V, Campell R. The effect of temperature and salinity on the toxicity of heavy metals to marine and estuarine invertebrates. *Oceanographic Marine Biology Annual Reviews* 1986;24:481-520.
 80. Depledge M. Interactions between heavy metals and physiological processes in estuarine invertebrates. In: Chambers PL, Chambers CM (Ed). *Estuarine Ecotoxicology*. Japaga; 1990. p. 89-100.
 81. Hall Jr. LW, Anderson RD. The influence of salinity on the toxicity of various classes of chemicals to aquatic biota. *Crit Rev Toxicol* 1995;25:281-346.
<http://dx.doi.org/10.3109/10408449509021613>
 82. Wright DA. Trace metals and major ion interactions in aquatic animals. *Marine Poll Bull* 1995;31:8-18.
[http://dx.doi.org/10.1016/0025-326X\(95\)00036-M](http://dx.doi.org/10.1016/0025-326X(95)00036-M)
 83. Lee B-G, Wallace WG, Luoma SN. Uptake and loss kinetics of Cd, Cr and Zn in the bivalves *Poamocorbula amurensis* and *Macoma balthica*: effects of size and salinity. *Mar Ecol Prog Ser* 1998;175:177-89.
<http://dx.doi.org/10.3354/meps175177>
 84. Ramachandran S, Swezey MJ, Hodson PV, Boudreau M, Courtenay SC, Lee K, et al. Influence of salinity and fish species on PAH uptake from dispersed crude oil. *Marine Poll Bull* 2006;52:1182-9.
<http://dx.doi.org/10.1016/j.marpolbul.2006.02.009>
 85. Casteel M, Sobsey M, Mueller J. Fecal contamination of agricultural soils before and after hurricane-associated flooding in North Carolina. *J Environ Sci Health A* 2006;41(2):173-84.
<http://dx.doi.org/10.1080/10934520500351884>
 86. Donald D, Hunter F, Sverko E, Hill B, Syrigiannis J. Mobilization of pesticides on an agricultural landscape flooded by a torrential storm. *Environ Toxicol Chem* 2005;24(1):2-10.
<http://dx.doi.org/10.1897/03-668.1>
 87. Boening DW. Ecological effects, transport, and fate of mercury: a general review. *Chemosphere* 2000;40(12):1335-51.
[http://dx.doi.org/10.1016/S0045-6535\(99\)00283-0](http://dx.doi.org/10.1016/S0045-6535(99)00283-0)
 88. United Nations Environment Programme. *Global mercury assessment*. Geneva: UNEP; 2002.
 89. Downs S, MacLeod C, Lester J. Mercury in precipitation and its relation to bioaccumulation in fish: a literature review. *Water Air Soil Poll* 1998;108:149-87.
 90. Booth S, Zeller D. Mercury, food webs, and marine mammals: implications of diet and climate change for human health. *Environ Health Perspect* 2005;113(5):521-6.
<http://dx.doi.org/10.1289/ehp.7603>
 91. Rodriguez M, Serodes J. Spatial and temporal evolution of trihalomethanes in three water distribution systems. *Water Res* 2001;35:1572-86.
[http://dx.doi.org/10.1016/S0043-1354\(00\)00403-6](http://dx.doi.org/10.1016/S0043-1354(00)00403-6)
 92. Zhang X, Echigo S, Lei H, Smith M, Minear R, Talley J. Effects of temperature and chemical addition on the formation of bromoorganic DBPs during ozonation. *Water Res* 2005;39:423-35.
<http://dx.doi.org/10.1016/j.watres.2004.10.007>
 93. Proia L, Vilches C, Boninneau C, Kantiani L, Farré M, Romani AM, et al. Drought episodically modulates the response of river biofilms to triclosan. *Aquatic Toxicol* 2012 (in press).
<http://dx.doi.org/10.1016/j.aquatox.2012.01.006>
 94. Guys CM, O'Keeffe J. Simple words and fuzzy zones: early directions for temporary river research in South Africa. *Environ Manage* 1997;21:517-31.
 95. Larned ST, Datry T, Arscott DB, Tockner K. Emerging concepts in temporary-river ecology. *Freshwater Biol* 2010;55(4):717-38.
<http://dx.doi.org/10.1111/j.1365-2427.2009.02322.x>
 96. Zoppini A, Amalfitano S, Fazi S, Puddu A. Dynamics of a benthic microbial community in a riverine environment subject to hydrological fluctuations (Mulargia River, Italy). *Hydrobiologia* 2010;657:37-51.
<http://dx.doi.org/10.1007/s10750-010-0199-6>
 97. Zoppini A, Marxsen J. Importance of extracellular enzymes for biogeochemical processes in temporary river sediments during fluctuating dry-wet conditions. In: Shukla G, Varma A (Ed.). *Soil enzymology*. Berlin Heidelberg: Springer; 2011. p. 103-17.
http://dx.doi.org/10.1007/978-3-642-14225-3_6

98. Ademollo N, Capri S, Patrolecco L, Puddu A, Polesello S, Rusconi M, *et al.* Fate and monitoring of hazardous substances in temporary rivers. *TrAC Trends in Analytical Chemistry* 2011;30(8):1222-32.
<http://dx.doi.org/10.1016/j.trac.2011.05.002>
99. Manganelli M, Scardala S, Stefanelli M, Palazzo F, Funari E, Vichi S, *et al.* Emerging Health Issues Of Cyanobacterial Blooms. *Ann Ist Super Sanità* 2012;48(4):415-28.
100. Scavia D, Field JC, Boesch DF, Buddemeier R, Cayan DR, Burkett V, *et al.* Climate change impacts on US coastal and marine ecosystems. *Estuaries* 2002;25:149-64.
101. Moore MV, Pace ML, Mather JR, Murdoch PS, Howarth RW, Folt CL, *et al.* Potential effects of climate change on freshwater ecosystems of the new england/mid-atlantic region. *Hydrological Processes* 1997;11(8):925-47.
[http://dx.doi.org/10.1002/\(sici\)1099-1085\(19970630\)11:8<925::aid-hyp512>3.0.co;2-x](http://dx.doi.org/10.1002/(sici)1099-1085(19970630)11:8<925::aid-hyp512>3.0.co;2-x)
102. Robson BJ, Hamilton DP. Summer flow event induces a cyanobacterial bloom in a seasonal Western Australian estuary. *Marine Freshwater Res* 2003;54(2):139-51.
<http://dx.doi.org/10.1071/MF02090>
103. Tonk L, Bosch K, Visser PM, Huisman J. Salt tolerance of the harmful cyanobacterium *Microcystis aeruginosa*. *Aquat Microb Ecol* 2007;46(2):117-23.
<http://dx.doi.org/10.3354/ame046117>
104. Ross C, Santiago-Vázquez L, Paul V. Toxin release in response to oxidative stress and programmed cell death in the cyanobacterium *Microcystis aeruginosa*. *Aquatic Toxicol* 2006;78(1):66-73.
<http://dx.doi.org/10.1016/j.aquatox.2006.02.007>
105. Jewel M, Affan M, Khan S. Fish mortality due to cyanobacterial bloom in an aquaculture pond in Bangladesh. *Pakistan J Biological Sci* 2003;6(12):1046-50.
106. Elmetri I, Bell PRF. Effects of phosphorus on the growth and nitrogen fixation rates of *Lynghya majuscula*: implications for management in Moreton Bay, Queensland. *Mar Ecol Prog Ser* 2004;281:27-35.
<http://dx.doi.org/10.3354/meps281027>
107. Albert S, O'Neil JM, Udy JW, Ahern KS, O'Sullivan CM, Dennison WC. Blooms of the cyanobacterium *Lynghya majuscula* in coastal Queensland, Australia: disparate sites, common factors. *Marine Poll Bull* 2005;51(1-4):428-37.
<http://dx.doi.org/10.1016/j.marpolbul.2004.10.016>
108. Jeppesen E, Kronvang B, Meerhoff M, Søndergaard M, Hansen KM, Andersen HE, *et al.* Climate change effects on runoff, catchment phosphorus loading and lake ecological state, and potential adaptations. *J Environ Quality* 2009;38(5):1930-41.
<http://dx.doi.org/10.2134/jeq2008.0113>
109. Paerl HW, Huisman J. Climate change: a catalyst for global expansion of harmful cyanobacterial blooms. *Environ Microbiol Rep* 2009;1(1):27-37.
<http://dx.doi.org/10.1111/j.1758-2229.2008.00004.x>
110. Robarts RD, Zohary T. Temperature effects on photosynthetic capacity, respiration, and growth rates of bloom-forming cyanobacteria. *New Zealand J Marine Freshwater Res* 1987;21(3):391-9.
<http://dx.doi.org/10.1080/00288330.1987.9516235>
111. Coles JF, Jones RC. Effect of temperature on photosynthesis-light response and growth of four phytoplankton species isolated from a tidal freshwater river. *J Phycol* 2000;36(1):7-16.
<http://dx.doi.org/10.1046/j.1529-8817.2000.98219.x>
112. Paerl HW, Xu H, McCarthy MJ, Zhu G, Qin B, Li Y, *et al.* Controlling harmful cyanobacterial blooms in a hyper-eutrophic lake (Lake Taihu, China): The need for a dual nutrient (N & P) management strategy. *Water Res* 2011;45(5):1973-83.
<http://dx.doi.org/10.1016/j.watres.2010.09.018>
113. Nalewajko C, Murphy TP. Effects of temperature, and availability of nitrogen and phosphorus on the abundance of *Anabaena* and *Microcystis* in Lake Biwa, Japan: an experimental approach. *Limnology* 2001;2:45-8.
<http://dx.doi.org/10.1007/s102010170015>
114. Elliott J, Jones I, Thackeray S. Testing the sensitivity of phytoplankton communities to changes in water temperature and nutrient load, in a temperate lake. *Hydrobiologia* 2006;559(1):401-11.
<http://dx.doi.org/10.1007/s10750-005-1233-y>
115. Chu Z, Jin X, Iwami N, Inamori Y. The effect of temperature on growth characteristics and competitions of *Microcystis aeruginosa* and *Oscillatoria mougeotii* in a shallow, eutrophic lake simulator system. *Hydrobiologia* 2007;581:217-23.
<http://dx.doi.org/10.1007/s10750-006-0506-4>
116. Peeters F, Straile D, Lorke A, Livingstone DM. Earlier onset of the spring phytoplankton bloom in lakes of the temperate zone in a warmer climate. *Global Change Biol* 2007;13(9):1898-909.
<http://dx.doi.org/10.1111/j.1365-2486.2007.01412.x>
117. Briand JF, Leboulanger C, Humbert JF, Bernard C, Dufour P. *Cylindrospermopsis raciborskii* (cyanobacteria) invasion at mid-latitudes: selection, wide physiological tolerance, or global warming? *JPhycol* 2004;40:231-8.
<http://dx.doi.org/10.1111/j.1529-8817.2004.03118.x>
118. Haande S, Rohrlack T, Ballot A, Røberg K, Skullberg R, Beck M, *et al.* Genetic characterisation of *Cylindrospermopsis raciborskii* (Nostocales, Cyanobacteria) isolates from Africa and Europe. *Harmful Algae* 2008;7(5):692-701.
<http://dx.doi.org/10.1016/j.hal.2008.02.010>
119. Saker ML, Griffiths DJ. The effect of temperature on growth and cylindrospermopsin content of seven isolates of *Cylindrospermopsis raciborskii* (Nostocales, Cyanophyceae) from water bodies in northern Australia. *Phycologia* 2000;39(4):349-54.
<http://dx.doi.org/10.2216/i0031-8884-39-4-349.1>
120. Lagos N, Onodera H, Zagatto P, Andrinolo D, Azevedo S, Oshima Y. The first evidence of paralytic shellfish toxins in the fresh water cyanobacterium *Cylindrospermopsis raciborskii*, isolated from Brazil. *Toxicon* 1999;37:1359-73.
[http://dx.doi.org/10.1016/S0041-0101\(99\)00080-X](http://dx.doi.org/10.1016/S0041-0101(99)00080-X)
121. Fastner J, Heinze R, Humpage AR, Mischke U, Eaglesham GK, Chorus I. Cylindrospermopsin occurrence in two German lakes and preliminary assessment of toxicity and toxin production of *Cylindrospermopsis raciborskii* (Cyanobacteria) isolates. *Toxicon* 2003;42(3):313-21.
[http://dx.doi.org/10.1016/S0041-0101\(03\)00150-8](http://dx.doi.org/10.1016/S0041-0101(03)00150-8)
122. Staal M, Meysman FJ, Stal LJ. Temperature excludes N₂-fixing heterocystous cyanobacteria in the tropical oceans. *Nature* 2003;425(6957):504-7.
<http://dx.doi.org/10.1038/nature01999>
123. Breitbarth E, Oschlies A, LaRoche J. Physiological constraints on the global distribution of *Trichodesmium* – effect of temperature on diazotrophy. *Biogeosciences* 2007;4:53-61.
<http://dx.doi.org/10.5194/bg-4-53-2007>
124. Huisman J, Sharples J, Stroom JM, Visser PM, Kardinaal WEA, Verspagen JMH, *et al.* Changes in turbulent mixing shift competition for light between phytoplankton species. *Ecology* 2004;85(11):2960-70.
<http://dx.doi.org/10.1890/03-0763>
125. Shatwell T, Köhler J, Nicklisch A. Warming promotes cold-adapted phytoplankton in temperate lakes and opens a loophole for *Oscillatoriales* in spring. *Global Change Biol* 2008;14(9):2194-200.
<http://dx.doi.org/10.1111/j.1365-2486.2008.01630.x>

126. Suikkanen S, Laamanen M, Huttunen M. Long-term changes in summer phytoplankton communities of the open northern Baltic Sea. *Estuarine, Coastal and Shelf Science* 2007;71(3-4):580-92.
<http://dx.doi.org/10.1016/j.ecss.2006.09.004>
127. Jacquet S, Briand JF, Leboulanger C, Avois-Jacquet C, Oberhaus L, Tassin B, et al. The proliferation of the toxic cyanobacterium *Planktothrix rubescens* following restoration of the largest natural French lake (Lac du Bourget). *Harmful Algae* 2005;4(4):651-72.
<http://dx.doi.org/10.1016/j.hal.2003.12.006>
128. Briand E, Yepremian C, Humbert JF, Quiblier C. Competition between microcystin- and non-microcystin-producing *Planktothrix agardhii* (cyanobacteria) strains under different environmental conditions. *Environ Microbiol* 2008;10(12):3337-48.
<http://dx.doi.org/10.1111/j.1462-2920.2008.01730.x>
129. Davis TW, Berry DL, Boyer GL, Gobler CJ. The effects of temperature and nutrients on the growth and dynamics of toxic and non-toxic strains of *Microcystis* during cyanobacteria blooms. *Harmful Algae* 2009;8(5):715-25.
<http://dx.doi.org/10.1016/j.hal.2009.02.004>
130. Oberholster P, Botha A-M, Myburgh J. Linking climate change and progressive eutrophication to incidents of clustered animal mortalities in different geographical regions of South Africa. *African J Biotechnol* 2009; 8(21):5825-32.
131. Sivonen K. Effects of light, temperature, nitrate, orthophosphate, and bacteria on growth of and hepatotoxin production by *Oscillatoria agardhii* strains. *Appl Environ Microbiol* 1990;56(9):2658-66.
132. Van der Westhuizen AJ, Eloff JN. Effect of temperature and light on the toxicity and growth of the blue-green alga *Microcystis aeruginosa* (UV-006). *Planta* 1985;163(1):55-9.
<http://dx.doi.org/10.1007/bf00395897>
133. Preußel K, Wessel G, Fastner J, Chorus I. Response of cylindrospermopsin production and release in *Aphanizomenon flos-aquae* (Cyanobacteria) to varying light and temperature conditions. *Harmful Algae* 2009;8(5):645-50.
<http://dx.doi.org/10.1016/j.hal.2008.10.009>
134. Dias E, Pereira P, Franca S. Production of paralytic shellfish toxins by *Aphanizomenon* sp. LMECYA 31 (cyanobacteria). *J Phycol* 2002;38(4):705-12.
<http://dx.doi.org/10.1046/j.1529-8817.2002.01146.x>
135. Rapala J, Sivonen K, Lyra C, Niemela S. Variation of microcystins, cyanobacterial hepatotoxins, in *Anabaena* spp. as a function of growth stimuli. *Appl Environ Microbiol* 1997; 63(6):2206-12.
136. Bowling LC, Baker PD. Major cyanobacterial bloom in the Barwon-Darling River, Australia, in 1991, and underlying limnological conditions. *Mar Freshwater Res* 1996;47:643-57.
<http://dx.doi.org/10.1071/MF9960643>
137. Walker SR, Lund JC, Schumacher DG, Brakhage PA, McManus BC, Miller JD, et al. Nebraska experience. In: Hudnell HK (Ed.). *Cyanobacterial harmful algal blooms: state of the science and research needs*. Springer New York; 2008. p. 139-52.
138. European Union. Directive 2007/60/EC, on the assessment and management of flood risks. *Official Journal of the European Union* L 288 6/11/2007.