

No physical health without mental health: lessons unlearned?

Kavitha Kolappa,^a David C Henderson^a & Sandeep P Kishore^b

Dr Brock Chisholm, the first Director-General of the World Health Organization (WHO), was a psychiatrist and shepherded the notion that mental and physical health were intimately linked. He famously stated that “without mental health there can be no true physical health”.¹ Half a century later, we have strong evidence elucidating the bidirectional relationship between mental illnesses – specifically depression and anxiety – and physical health outcomes. However, policy continues to lag behind the evidence in this regard, as demonstrated by our global noncommunicable disease response.

Over a decade ago, the World Health Assembly adopted a global strategy for the prevention and control of noncommunicable disease. At the time, these were limited to the following four illness types: cardiovascular disease, diabetes, respiratory illness and cancers. Such a categorization would set a precedent for the exclusion of mental illnesses from all future WHO discussions on noncommunicable diseases. It is not surprising then, that in the *2008–2013 action plan for the global strategy for the prevention and control of noncommunicable diseases* mental illnesses were relegated to a footnote, with the justification that they do not share risk factors with the other four types of illnesses.²

We take issue with this viewpoint, as mental illnesses are themselves risk factors that affect the incidence and prognosis of diseases traditionally classified as “noncommunicable”. Patients with type II diabetes mellitus, for example, are twice as likely to experience depression as the general population,³ and those patients with diabetes who are depressed have greater difficulty with self-care.⁴ Patients suffering from mental illness are twice as likely to

smoke cigarettes as other people, and in patients with chronic obstructive pulmonary disease mental illness is linked to poorer clinical outcomes.^{5,6} Up to 50% of cancer patients suffer from a mental illness, especially depression and anxiety,⁷ and treating symptoms of depression in cancer patients may improve survival time.⁸ Similarly, in patients who are depressed, the risk of having a heart attack is more than twice as high as in the general population;⁹ further, depression increases the risk of death in patients with cardiac disease.¹⁰ Moreover, treating the symptoms of depression after a heart attack has been shown to lower both mortality and re-hospitalization rates.¹¹ In light of this evidence, how can we possibly address the burgeoning epidemic of noncommunicable diseases without tackling co-morbid mental illnesses?

Mental illnesses were declared a regional priority in Africa during the WHO African Region Ministerial Consultation on Noncommunicable Diseases, held in Brazzaville, Congo, in April 2011. Later that month the WHO's African Member States and India reiterated this priority at the first Global Ministerial Conference on Healthy Lifestyles and Noncommunicable Disease Control, held in Moscow, Russia.¹² As a result, mental illnesses were featured prominently in the preambles of the Moscow Declaration, as well as in the political declaration issued by the United Nations General Assembly at the high-level meeting on noncommunicable diseases held in New York City in September 2011.¹³ Despite this progress, however, mental illnesses received no mention at all in the resolution on noncommunicable diseases that WHO's Member States adopted during the 130th session of WHO's Executive Board.¹⁴ Mental

illnesses were also omitted from WHO's proposed monitoring framework, indicators and voluntary targets for the prevention and control of noncommunicable diseases, which was released in November 2012.¹⁵

The *2008–2013 action plan for the global strategy for the prevention and control of noncommunicable diseases* will be revised over the coming year, and the WHO's Executive Board and World Health Assembly are preparing their deliberations for 2013. During this critical time we urge Member States to recognize the importance of co-morbid mental illnesses as amplifiers of the burden of other noncommunicable diseases. To this end, we call on Member States to assess and monitor co-morbid mental illnesses in primary care settings, prioritize the training of professionals in mental health care, and, critically, incorporate mental health interventions within chronic disease programs as part of a vigorous global response to noncommunicable diseases. We now know that addressing mental illnesses in primary care settings will delay progression, improve survival outcomes, and reduce the health care costs of other noncommunicable diseases. The time has now come to do away with the artificial divisions between mental and physical health, as WHO's first Director-General championed so many decades ago. ■

References

Available at: <http://www.who.int/bulletin/volumes/91/1/12-115063>

^a Massachusetts General Hospital, Boston, United States of America (USA).

^b Young Professionals Chronic Disease Network, 1230 York Avenue, Box 292, New York, NY 10065, USA.

Correspondence to Sandeep P Kishore (e-mail: sunny.kishore@gmail.com).

References

1. *Outline for a Study Group on World Health and the Survival of the Human Race. Material drawn from articles and speeches by Brock Chisholm.* Geneva: World Health Organization; 1954. Available from: whqlibdoc.who.int/hist/.../ChisholmBrock_1953_Compilation.pdf [accessed 6 December 2012].
2. *2008–2013 action plan for the global strategy for the prevention and control of noncommunicable diseases.* Geneva: World Health Organization; 2009. Available from: <http://www.who.int/nmh/publications/9789241597418/en/> [accessed 30 November 2012].
3. Cosgrove MP, Sargeant LA, Griffin SJ. Does depression increase the risk of developing type 2 diabetes? *Occup Med (Lond)* 2008;58:7–14. doi:10.1093/occmed/kqm105 PMID:17965449
4. Gonzalez JS, Safren SA, Delahanty LM, Cagliero E, Wexler DJ, Meigs JB et al. Symptoms of depression prospectively predict poorer self-care in patients with Type 2 diabetes. *Diabet Med* 2008;25:1102–7. doi:10.1111/j.1464-5491.2008.02535.x PMID:19183315
5. Lasser K, Boyd JW, Woolhandler S, Himmelstein DU, McCormick D, Bor DH. Smoking and mental illness: a population-based prevalence study. *JAMA* 2000;284:2606–10. doi:10.1001/jama.284.20.2606 PMID:11086367
6. Felker B, Bush KR, Harel O, Shofer JB, Shores MM, Au DH. Added burden of mental disorders on health status among patients with chronic obstructive pulmonary disease. *Prim Care Companion J Clin Psychiatry* 2010;12:pii: PCC.09m00858. PMID:21085558
7. Massie MJ. Prevalence of depression in patients with cancer. *J Natl Cancer Inst Monogr* 2004;32:57–71. doi:10.1093/jncimonographs/lgh014 PMID:15263042
8. Giese-Davis J, Collie K, Rancourt KM, Neri E, Kraemer HC, Spiegel D. Decrease in depression symptoms is associated with longer survival in patients with metastatic breast cancer: a secondary analysis. *J Clin Oncol* 2011;29:413–20. doi:10.1200/JCO.2010.28.4455 PMID:21149651
9. Rugulies R. Depression as a predictor for coronary heart disease. a review and meta-analysis. *Am J Prev Med* 2002;23:51–61. doi:10.1016/S0749-3797(02)00439-7 PMID:12093424
10. Whang W, Shimbo D, Kronish IM, Duvall WL, Julien H, Iyer P et al. Depressive symptoms and all-cause mortality in unstable angina pectoris (from the Coronary Psychosocial Evaluation Studies [COPEs]). *Am J Cardiol* 2010;106:1104–7. doi:10.1016/j.amjcard.2010.06.015 PMID:20920647
11. Mazza M, Lotrionte M, Biondi-Zoccai G, Abbate A, Sheiban I, Romagnoli E. Selective serotonin reuptake inhibitors provide significant lower re-hospitalization rates in patients recovering from acute coronary syndromes: evidence from a meta-analysis. *J Psychopharmacol* 2010;24:1785–92. doi:10.1177/0269881109348176 PMID:19965939
12. Dhar A. At India's insistence, mental health included as non-communicable disease. *The Hindu*. 4 May 2011. Available from: <http://www.thehindu.com/health/article1991279.ece> [accessed 3 December 2012].
13. Resolution A/RES.66/2. Political Declaration of the High-level Meeting of the General Assembly on the Prevention and Control of Non-communicable Diseases. In: *Sixty-sixth General Assembly of the United Nations, New York, 19–20 September 2011* [Internet]. New York: UNGA; 2012. Available from: <http://www.un.org/en/sc/documents/resolutions/> [accessed 3 December 2012].
14. Resolution EB130.R7. Prevention and control of non-communicable diseases: Follow-up to the High-level Meeting of the United Nations General Assembly on Prevention and Control of Non-communicable Diseases. In: *130th session of the Executive Board of the World Health Organization, Geneva, 16–23 January 2012* [Internet]. Geneva: WHO; 2012. Available from: apps.who.int/gb/ebwha/pdf_files/EB130/B130_6-en.pdf [accessed 3 December 2012].
15. *Report of the formal meeting of Member States to conclude the work on the comprehensive global monitoring framework, including indicators, and a set of voluntary global targets for the prevention and control of noncommunicable diseases.* Geneva: World Health Organization; 2012. Available from: <http://apps.who.int/gb/ncds/> [accessed 3 December 2012].