

physical illness, for example, can impair the psychological health of their female partners; when men are sick, injured or die, households and female partners suffer a loss of income.²³ Closing the men's health gap can benefit men, women and their children. ■

Acknowledgements:

We thank Sarah Hawkes, Reader in Global Health at University College London's Institute of Global Health, for her valuable comments. Tim Shand thanks Dr Hawkes for being his PhD supervisor at University College London.

Competing interests: PB reports an honorarium from Eli Lilly and Company to

conduct a training session on men's health, two honoraria from Sanofi Pasteur MSD (one for a presentation in 2012 to an advisory group on HPV vaccination for boys and one for contributing in 2013 to a study looking at developing a model health technology assessment for HPV vaccination with boys as the case study), and an honorarium from Nicholas Hall and Company. SLD reports no relevant competing interests. ST declares that he has no relevant competing interests. IB reports funding for travel expenses from the British Medical Association, Karolinska Institute, Ulster University, BMS, the Northern Ireland Health Service, the European Cancer Organisation, the European Commission, and Nicholas Hall,

and funding for project work from GSK, Pfizer, Astellas, SCA Sweden, Amgen, and Proctor & Gamble. GY declares that E2Pi has received funding from the Bill & Melinda Gates Foundation, the Global Fund, the Clinton Health Access Initiative, UNITAID, the Harvard Global Health Institute, the Disease Control Priorities Network, the UK Department for International Development, and the Norwegian Agency for Development Cooperation. TS reports that Sonke Gender Justice receives funding from the Swedish International Development Cooperation Agency, the Norwegian Agency for Development Cooperation, UNFPA, and the US Agency for International Development.

References

- Wang H, Dwyer-Lindgren L, Lofgren KT, Rajaratnam JK, Marcus JR, Levin-Rector A, et al. Age-specific and sex-specific mortality in 187 countries, 1970–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2071–94.
- Jamison DT, Summers LH, Alleyne G, Arrow KJ, Berkley S, Binagwaho A, et al. Global Health 2035: a world converging within a generation. *Lancet*. 2013;382:1898–955.
- UCL Institute of Health Equity [Internet]. Review of social determinants and the health divide in the WHO European Region: final report. Copenhagen: World Health Organization, Regional Office for Europe; 2013. Available from: <http://www.instituteoftheequity.org/projects/who-european-review> [cited 2014 Feb 21].
- Lim S, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2224–60.
- Hinote BP, Webber GR. Drinking toward manhood: masculinity and alcohol in the former USSR. *Men Masc*. 2012;15:292–310.
- Hawkes S, Buse K. Gender and global health: evidence, policy, and inconvenient truths. *Lancet*. 2013;381:1783–7.
- Hippisley-Cox J, Vinogradova Y. Trends in consultation rates in general practice 1995/1996 to 2008/2009: analysis of the QResearch® database. Leeds: Health and Social Care Information Centre, 2009.
- Sakalauskiene Ž, Vehkalahti MM, Murtomaa H, Mačičiuskienė V. Factors related to gender differences in toothbrushing among Lithuanian middle-aged university employees. *Medicina (Kaunas)*. 2011;47:180–6.
- Skovdal M, Campbell C, Madanhire C, Mupambireyi Z, Nyamukapa C, Gregson S. Masculinity as a barrier to men's use of HIV services in Zimbabwe. *Global Health*. 2011;7:13.
- Cornell M, McIntyre J, Myer L. Men and antiretroviral therapy in Africa: our blind spot. *Trop Med Int Health*. 2011;16:828–9.
- European Commission [Internet]. The state of men's health in Europe report. Brussels: European Union; 2011. Available from: http://ec.europa.eu/health/population_groups/docs/men_health_report_en.pdf [cited 2014 June 6].
- Centers for Disease Control and Prevention. Workers Memorial Day – April 28, 2012. *MMWR Morb Mortal Wkly Rep*. 2012;61:281.
- McKinlay E, Kljakovic M, McBain L. New Zealand men's health care: are we meeting the needs of men in general practice? *J Primary Health Care* 2009;1(4):302–10.
- Barker G, Ricardo C, Nascimento M, Olukoya A, Santos C. Questioning gender norms with men to improve health outcomes: evidence of impact. *Glob Public Health*. 2010;5:539–53.
- Juel K, Christensen K. Are men seeking medical advice too late? Contacts to general practitioners and hospital admissions in Denmark 2005. *J Public Health (Oxf)*. 2008;30:111–3.
- Brott A, Dougherty A, Williams ST, Matope JH, Fadich A, Taddelle M. The economic burden shouldered by public and private entities as a consequence of health disparities between men and women. *Am J Mens Health*. 2011;5:528–39.
- White A, McKee M, Richardson N, de Visser R, Madsen SA, de Sousa BC. Europe's men need their own health strategy. *BMJ*. 2011;343:d739.
- Hunt K, Wyke S, Gray CM, Anderson AS, Brady A, Bunn C, et al. A gender-sensitised weight loss and healthy living programme for overweight and obese men delivered by Scottish Premier League football clubs (FFIT): a pragmatic randomised controlled trial. *Lancet*. 2014;21 Jan.
- Brugha RF, Kevany JP, Swan AV. An investigation of the role of fathers in immunization uptake. *Int J Epidemiol*. 1996;25:840–5.
- Lindegren ML, Kennedy CE, Bain-Brickley D, Azman H, Creanga AA, Butler LM, et al. Integration of HIV/AIDS services with maternal, neonatal and child health, nutrition, and family planning services. *Cochrane Database Syst Rev*. 2012;9:CD010119.
- Dworkin SL, Treves-Kagan S, Lippman SA. Gender-transformative interventions to reduce HIV risks and violence with heterosexually active men: a review of the global evidence. *AIDS Behav*. 2013;17:2845–63.
- World Health Organization, Regional Office for Europe [Internet]. Men's Health. Geneva: WHO; 2014. Available from: <http://www.euro.who.int/en/health-topics/health-determinants/gender/activities/mens-health> [cited 2014 Feb 21].
- Hagedoorn M, Sanderman R, Ranchor AV, Brilman EI, Kempen GJ, Ormel J. Chronic disease in elderly couples: are women more responsive to their spouses' health condition than men? *J Psychosom Res*. 2001;51:693–6.

Corrigendum

In Volume 92, Issue 7, July 2014, page 533, the 27th author should be spelt "Susan Jack".

In Volume 92, Issue 5, May 2014, page 340, should have:

- addition of an affiliation "d" as "Chinese Center for Disease Control and Prevention, Beijing, China."
- correction of affiliation for Xiaofeng Liang and Weizhong Yang to "d"
- correction of affiliation "b" to "National Institute for Viral Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing, China."

In Volume 88, Issue 4, April 2010, page 279, the second sentence of the second paragraph should read:

ودعا الباحثون جميع السكان في عمر اكبر من أو يساوي 15 سنة لإجراء فحص السعال وفحص الأشعة السينية عليهم.