

COVID-19, social vulnerability and mental health of LGBTQIA+ populations

COVID-19, vulnerabilidade social e saúde mental das populações LGBTQIA+

COVID-19, vulnerabilidad social y salud mental de las poblaciones LGBTQIA+

Geovani Bordiano ¹
Suzana Pacheco Liberal ¹
Giovanni Marcos Lovisi ¹
Lucia Abelha ¹

doi: 10.1590/0102-311X00287220

A significant aspect observed during the evolution of the COVID-19 pandemic is the relationship between social vulnerability and higher prevalence and lethality of the virus ¹. The contrasts of this mortality between the poorest and richest show how social inequalities directly affect the Brazilian population's health and make certain social groups more vulnerable ¹.

In addition to the damages to health already known, the pandemic can be considered a traumatic event and thus generate serious consequences for the mental health of the population as a whole, namely: insomnia, depression, fear and anger reactions, substance abuse, acute reactions to stress, and posttraumatic stress disorder ². Thus, the pandemic can be understood as a catalyst for mental illness, especially for the most vulnerable populations.

Sexual and gender minorities, i.e., lesbian, gay, bisexual, transvestite, transsexual and transgender, queers, intersex, asexual, and other (LGBTQIA+) have suffered, and still suffer, a serious process of social exclusion throughout the history of contemporary societies ³. One of the consequences of this fact is the worsening of the mental health status of these groups.

Aspects such as low schooling level, low socioeconomic status, violence, and difficulties in accessing health services are closely related to social minorities and, at the same time, they have been considered risk factors for common mental disorders in the general population ^{4,5}. Furthermore, other risk factors for mental health are specific to these marginalized groups: prejudice (or LGBTQIA+phobia) ⁶, lack of institutionalized protection, family rejection, and bullying ⁷. This whole social conjuncture cooperates for the fragility and vulnerability of these minorities.

Therefore, national and international studies ^{8,9,10,11,12,13,14} reveal that sexual and gender minorities – when compared to the non-LGBTQIA+ population – present higher prevalence of depression and anxiety, higher risk for suicide and they make a more intense use of psychoactive substances.

Pandemic, mental health and LGBTQIA+ populations

In a recent national survey ¹³, conducted with about 9,000 participants, which aimed to investigate the greatest effects of the pandemic for the LGBTQIA+ population, 42.72% of respondents indicated mental health as the main effect experienced during this period, followed by “new rules of conviviality”, “loneliness”, “family life”, and “lack of employment and money”.

¹ Instituto de Estudos em Saúde Coletiva, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brasil.

Correspondence

G. Bordiano
Instituto de Estudos em Saúde Coletiva, Universidade Federal do Rio de Janeiro, Av. Horácio Macedo s/n, Rio de Janeiro, RJ 21941-598, Brasil.
giovannibordiano@gmail.com



Among the reasons that make LGBTQIA+ more vulnerable at times like the current one, we can mention the high rates of violence to which they are subjected daily. Despite the lack of data, figures collected by the Institute for Applied Economic Research (IPEA) in the *Atlas of Violence* ¹⁵ show a 127% increase in reports of LGBTQIA+ homicide in 2018 in Brazil. Moreover, one of the most important health measures recommended by health agencies has been social distancing that, in the case of LGBTQIA+, may mean remaining confined to potential aggressors, often their own families or intimate partners ^{11,12,13,16}.

The increase in domestic violence due to social distancing has been brought to the discussions ¹⁷, but generally with an exclusive focus on heterosexual cisgender women. However, it is noteworthy that among the homicides by LGBTQIA+phobia in Brazil, the victim's residence is the second place of highest occurrence of murders ¹⁸, with transsexual, transgender, and transvestite people being preferred targets of this type of violence, followed by lesbians and then gay men ¹⁶.

Socioeconomic vulnerability is also a relevant aspect, permeating income and labor bonds. It is known that the pandemic has caused job losses and precarious employment ties – that were already fragile – which may aggravate the socioeconomic situation of these populations ^{11,12}. An North American study indicates that the largest portion of this population in the country is part of the informal market and also has occupations potentially more exposed to SARS-CoV-2, such as work in bars and restaurants, in addition to having lower income and lower conditions of access to health ¹⁹. In Brazil, when considering only the populations of transgender, transsexual and transvestite, vulnerability is even greater given the specificities of these groups ¹². A study conducted with 672 transsexuals in the State of São Paulo revealed that, from 2014 to 2015, only 16.7% were in the formal market ²⁰ and, still, many worked as sex workers ²¹. Regarding the homosexual and bisexual population, an international systematic review ²² indicates that gay and bisexual men receive less than their heterosexual colleagues. Furthermore, the unemployment rate in this population, found by a survey ¹³, is 21.6% in Brazil and, during the pandemic, 44.3% suffered a total shutdown of their activities.

Social life is another significant aspect placed on frailty for LGBTQIA+ during the pandemic. As they are sometimes rejected in the family nucleus, a significant mechanism of resistance and survival of these groups lies in the bond they establish with their communities, so that, considering the social distancing measures, these individuals experience a serious impact because of the impossibility of a more active and face-to-face contact with their affections and familiar places, which leads to an experience of isolation and loneliness ^{11,12,13}.

Considering so many aspects that compromise LGBTQIA+ individuals' mental health, it is indispensable that the authorities define care strategies for this moment.

The Brazilian Ministry of Health – in its booklet on suicide in the COVID-19 pandemic ²³ – lists some warning factors that deserve special attention, among them: preexisting mental health disorders, with emphasis on depression and anxiety; use of alcohol and/or other drugs; lack of social support; existence of hostility in the family environment; and suffering and concerns about one's own sexuality. In another booklet ²⁴ that deals with recommendations for public administrators, the Brazilian Ministry of Health indicates that one of the points that should receive attention is the identification of vulnerable groups from the psychosocial point of view, and it highlights several recommendations that can be put in force by public administrators during the pandemic to ensure access to mental health in this population. However, it is clear that despite its significance, the existence of legal instruments, guidelines and regulations, it is not enough to assure the population's access to their rights.

Often these groups are exposed to prejudice even in health services, which can lead to decreased demand and adherence to treatments ^{25,26}. Thus, it is observed that the heteronormativity present in several care fields produces a devaluation or non-recognition of these identities, contributing to the perpetuation of violence and discrimination that affect the conditions of access and quality of health care ²⁷. A study ²⁷ reports that the main difficulties in accessing health services for this public are related to the fear of suffering prejudices and the fear of revealing sexual orientation, thus, this information is often omitted in care.

In the Brazilian Unified National Health System (SUS), specialized mental health care is provided by the Psychosocial Care Network (RAPS), which is one of the main services provided in psychosocial care centers (CAPS). However, it must be emphasized that the RAPS faces significant obstacles, such as insufficient equipment in some Brazilian regions and municipalities ²⁹, which leads to long waiting

lines, and reference and counter-reference problems³⁰. It is also noteworthy, considering the great percentage of discrimination suffered in health services, demonstrated in studies³¹, the lack of qualification of health professionals to deal with the sufferings related to sexuality.

Considering these data, it is possible to say that the mental health of sexual and gender minorities lacks, not only specific attention in terms of emergency public health policies during the pandemic, but, above all, mobilization and concrete actions. However, this data attest a profound invisibility of the issue by the State and sometimes also by academia³², even during a time of extreme need. Moreover, it is urgent that such discussion reach social spaces and that the LGBTQIA+ population achieve more visibility. Also, that the current situation be a lesson for future moments.

Contributors

G. Bordiano contributed to the project design, analysis and interpretation of data, writing, critical review of the intellectual content, and approval of the final version of the article; and is responsible for all aspects of the work. S. P. Liberal and G. M. Lovisi contributed to the project design, analysis and interpretation of data, writing, critical review of the intellectual content, and approval of the final version of the article. L. Abelha contributed to the critical review of the intellectual content and approval of the final version of the article.

Additional informations

ORCID: Geovani Bordiano (0000-0002-3285-0689); Suzana Pacheco Liberal (0000-0001-9890-6895); Giovanni Marcos Lovisi (0000-0003-0521-0202); Lucia Abelha (0000-0002-2409-9872).

References

1. Fundação Oswaldo Cruz. Análise da frequência, incidência, mortalidade e letalidade por COVID-19 em favelas cariocas. Boletim Socioepidemiológico da COVID-19 nas Favelas 2020; (1). https://portal.fiocruz.br/sites/portal.fiocruz.br/files/documentos/boletim_socioepidemiologicos_covid_nas_favelas_1.pdf.
2. Shigemura J, Ursano RJ, Morganstein JC, Kurosawa M, Benedek DM. Public responses to the novel 2019 coronavirus (2019-nCoV) in Japan: mental health consequences and target populations. *Psychiatry Clin Neurosci* 2020; 74:281-2.
3. Oliveira FAG, Carvalho HR, Jesus JG. LGBTI+ em tempos de pandemia da Covid-19. *Diversitates International Journal* 2020; 12:60-94.
4. Patel V, Kleinman A. Poverty and common mental disorders in developing countries. *Bull World Health Organ* 2003; 81:609-15.
5. Fundação Oswaldo Cruz. Saúde mental e atenção psicossocial na pandemia COVID-19: recomendações gerais. Rio de Janeiro: Fundação Oswaldo Cruz; 2020.
6. King M, Semlyen J, Tai SS, Killaspy H, Osborn D, Popelyuk D, et al. A systematic review of mental disorder, suicide, and deliberate self harm in lesbian, gay and bisexual people. *BMC Psychiatry* 2008; 8:70.
7. Shah R, Eshel N, McGlynn L. Lesbian, gay, bisexual, transgender, and queer/questioning students. In: Roberts LW, editor. *Student mental health: a guide for psychiatrists, psychologists, and leaders serving in higher education*. Washington DC: American Psychiatric Association Publishing; 2018. p. 416-9.
8. Di Giacomo E, Krausz M, Colmegna F, Aspesi F, Clerici M. Estimating the risk of attempted suicide among sexual minority youths. *JAMA Pediatr* 2018; 172:1145-52.
9. Teixeira-Filho FS, Rondini CA. Suicide thoughts and attempts of suicide in adolescents with hetero and homoerotic sexual practices. *Saúde Soc* 2012; 21:651-67.

10. Remy LS, Scherer J, Guimarães L, Surratt HL, Kurtz SP, Pechansky F, et al. Anxiety and depression symptoms in Brazilian sexual minority ecstasy and LSD users. *Trends Psychiatry Psychother* 2017; 39:239-46.
11. OutRight Action International. Vulnerability amplified: the impact of the COVID-19 pandemic on LGBTIQ people. New York: Out-Right Action International; 2020.
12. Carman M, Bourne A, Fairchild J. COVID-19 impacts for LGBTIQ communities and implications for services: a research briefing paper by Rainbow Health Victoria. Melbourne: Rainbow Health Victoria; 2020.
13. #VOTELGBT. Diagnóstico LGBT+ na pandemia. <https://www.votelgbt.org/pesquisas> (accessed on Sep/2020).
14. Francisco LCF, Barros AC, Pacheco M, Nardi AE, Alves V. Anxiety in sexual and gender minorities: an integrative review. *J Bras Psiquiatr* 2020; 69:48-56.
15. Cerqueira D, Bueno S, Lima RS, Neme C, Ferreira H, Alves PP, et al. Atlas de violência 2019. Brasília/Rio de Janeiro/São Paulo: Instituto de Pesquisa Econômica Aplicada/Fórum Brasileiro de Segurança Pública; 2019.
16. Pinto IV, Andrade SSA, Rodrigues LL, Santos MAS, Marinho MMA, Benício LA, et al. Perfil das notificações de violências em lésbicas, gays, bissexuais, travestis e transexuais registradas no Sistema de Informação de Agravos de Notificação, Brasil, 2015 a 2017. *Rev Bras Epidemiol* 2020; 23 Suppl 1:e200006.supl.1.
17. Vieira PR, Garcia LP, Maciel ELN. The increase in domestic violence during the social isolation: what does it reveals? *Rev Bras Epidemiol* 2020; 23:e200033.
18. Mendes WG, Silva CMFP. Homicide of lesbians, gays, bisexuals, travestis, transsexuals, and transgender people (LGBT) in Brazil: a spatial analysis. *Ciênc Saúde Colet* 2020; 25:1709-22.
19. Whittington C, Hadfield K, Calderón C. The lives and livelihoods of many in the LGBTQ community are at risk amidst COVID-19 crisis. Washington DC: Human Right Campaign Foundation; 2020.
20. Silva MA, Carla GL, Veras MASM. Work and health issues of the transgender population: factors associated with entering the labor market in the state of São Paulo, Brazil. *Ciênc Saúde Colet* 2020; 25:1723-34.
21. Reidel M. A pedagogia do salto alto: histórias de professoras transexuais e travestis na educação brasileira [Dissertação de Mestrado]. Porto Alegre: Universidade Federal do Rio Grande do Sul; 2013.
22. Ozeren E. Sexual orientation discrimination in the workplace: a systematic review of literature. *Procedia Soc Behav Sci* 2014; 109:1203-15.
23. Fundação Oswaldo Cruz. Saúde mental e atenção psicossocial na pandemia de COVID-19: suicídio na pandemia de COVID-19. Rio de Janeiro: Fundação Oswaldo Cruz; 2020.
24. Fundação Oswaldo Cruz. Saúde mental e atenção psicossocial na pandemia de COVID-19: recomendações para gestores. Rio de Janeiro: Fundação Oswaldo Cruz; 2020.
25. Baptiste-Roberts K, Oranuba E, Werts N, Edwards LV. Addressing health care disparities among sexual minorities. *Obstet Gynecol Clin North Am* 2017; 44:71-80.
26. Dahlhamer JM, Galinsky AM, Joestl SS, Ward BW. Barriers to health care among adults identifying as sexual minorities: a US national study. *Am J Public Health* 2016; 106:1116-22.
27. Ferreira BO, Bonan C. Opening the closets of access and quality: an integrative review on the health of lgbtt populations. *Ciênc Saúde Colet* 2020; 25:1765-78.
28. Lovisi GM, Abelha L, Schilling S, Muñoz RA. Prospects and challenges of mental health services in a major city in Latin America: Rio de Janeiro. In: Okkels N, Kristiansen CB, Munk-Jorgensen P, editors. *Mental health and illness in the city*. Singapore: Springer; 2017. p. 357-75.
29. Gama CAP, Guimarães DA, Coelho VAA, Carvalho RC, Campos CG, Fraga AMS. A implantação da rede de atenção psicossocial na Região Ampliada de Saúde Oeste de Minas Gerais – BR. *Cad Saúde Colet (Rio J.)* 2020; 28:278-87.
30. Quinderé PHD, Jorge MSB, Franco TB. Rede de atenção psicossocial: qual o lugar da saúde mental? *Physis (Rio J.)* 2014; 24:253-71.
31. McClain Z, Thomas R, Yehia BR. Sociocultural and systemic barriers to health for gender and sexual minority populations. In: Smalley KB, Warren JC, Barefoot, Nikki K, editors. *LGBT health: meeting the needs of gender and sexual minorities*. New York: Springer; 2018. p. 16-8.
32. Bezerra MV, Moreno CA, Prado NM, Santos AM. Política de saúde LGBT e sua invisibilidade nas publicações em saúde coletiva. *Saúde Debate* 2019; 43(n.spe 8):305-23.

Submitted on 01/Oct/2020

Final version resubmitted on em 25/Jan/2021

Approved on 12/Feb/2021