

Spatiotemporal Analysis of Lung Cancer Incidence and Case Fatality in Villa Clara Province, Cuba

Norma E. Batista MD and Oscar A. Antón MD

ABSTRACT

INTRODUCTION Cancer has historically been a main cause of death in Cuba, with lung cancer the number one cause of cancer death in both sexes. Cancer morbidity and mortality rates are the basic measures of cancer impact in the community. Cancer mortality has been one of the major applications of geographic analysis and has made important progress in recent decades thanks to access to mortality statistics and to development and availability of geographic information systems. Cuba does not have a strong tradition of etiologic research using spatial analysis. High levels of lung cancer morbidity and mortality in Villa Clara and growing interest in spatial analysis as an epidemiologic tool motivated this study.

OBJECTIVE To identify spatial and/or spatiotemporal clusters of lung cancer morbidity and case fatality in the province of Villa Clara, and to demonstrate the value of cluster analysis as an epidemiologic tool.

METHODS Descriptive observational study based on administrative data, using the technique of space-time scan statistics. The study

focused on new cases diagnosed in 2004 and case-fatality for those cases through 2009. Variables used were: cases diagnosed, deaths, date of diagnosis, date of death, municipality and Cartesian geocoding for each municipality.

RESULTS The study identified significant spatial and spatiotemporal clusters of greater than expected lung cancer incidence (municipalities of Encrucijada, Camajuaní, Cifuentes, Sagua la Grande, Caibarién and Santa Clara) and case fatality (Encrucijada, Camajuaní, Cifuentes, Sagua la Grande, Caibarién, Santa Clara, Placetas and Manicaragua).

CONCLUSIONS Although the results are not explanatory, the spatial and spatiotemporal patterns of excess lung cancer risk and case-fatality can support hypothesis generation for research and eventual interventions for targeted prevention and management.

KEYWORDS Pulmonary neoplasms, epidemiological methods, spatiotemporal analysis, cluster analysis, space-time clustering, incidence, case fatality, Cuba

INTRODUCTION

Cancer is a leading cause of death globally; each year 9 million people are diagnosed with cancer and approximately 5 million die. [1] This high morbidity and mortality, and frequently unclear etiology, make cancer a major health problem. Among the main factors associated with rising cancer rates are smoking, in both men and women, and population aging.[1,2] The latter also makes cancer an urgent social and economic challenge, with more people reaching the ages of highest cancer risk, due to economic and social development and health system improvements.[3,4]

In 2008, WHO reported Cuba's cancer mortality rate (age adjusted to the WHO world standard population) at 129 per 100,000 population, compared to 132 per 100,000 globally. Cuba ranked 103rd (highest to lowest) among the 193 countries included in the report, 6th in the Caribbean and 15th in the Americas (after Bolivia, Grenada, Peru, Uruguay, Jamaica, Antigua and Barbuda, Argentina, Paraguay, Honduras, Canada, Chile, Barbados, the United States and the Dominican Republic, in that order).[1] For over three decades, while cancer mortality trends varied, cancer was consistently the second cause of death for all ages in the Cuban population—surpassing cardiovascular disease to become first in 2011—and the first cause of years of potential life lost.[5,6]

In 2010, the central Cuban province of Villa Clara had a population of 800,300 with a population density of 95.1 inhabitants per km². The main industries were agriculture (sugar and tobacco), chemicals and metallurgy. The provincial cancer mortality rate (age adjusted to the 1981 Cuban census population) in 2009 was 201.3 per 100,000 population; this was above the national average (189.7 per 100,000 population), and surpassed the mortality rate from cardiovascular diseases (194.6 per 100,000 population), making Villa Clara one of the provinces with the highest cancer burden.[7]

Lung cancer is the most frequent cancer in Cuban men and the third most frequent in Cuban women; it is the first cause of cancer death in both sexes, both nationally and in Villa Clara where mortality rates in 2009 were 54.7 and 21.6 per 100,000 population for men and women, respectively.[7]

Air pollution is a serious environmental problem worldwide, regardless of level of economic development. It differs from other types of pollution since once contaminants are in the air, exposure cannot be easily avoided. Indoor air pollution (in houses, work centers, public places) presents many challenges; it is sometimes worse than outdoor air pollution.[8,9] Household air pollution is a major concern of researchers around the world, due to the prolonged and continuous exposure; a typical individual spends at least 50% of his/her time in the home. It has become the main source of exposure for children.[10,11]

Air contaminants are more common in the city than in the countryside, due to industrial waste emissions; motor vehicle exhaust; sulphur dioxide, one of the most important carcinogenic reducing agents; pavement tar particles; and radioactive particles from naturally occurring radon gas decay (the main cause of lung cancer in nonsmokers).[12] Some 75.8% of Villa Clara's population reside in urban settings.[7]

Although most authors acknowledge the existence of an urban factor in lung cancer incidence, the main culprit by far is smoking.[12–15] Tobacco use is responsible for high cancer incidence and mortality in Cuba, contributing to 15% of all deaths in 1995 and 18% in 2007, surpassing attributable mortality reported in the adult population globally (12%) and in countries such as Canada (16.6%), Paraguay (13.4%) and France (12%). Tobacco use caused 86% of the 1209 lung cancer deaths in Cuba in 2007. Seven out of every ten tobacco-associated cancer deaths were in

men, 64–65% of these from tracheobronchial–pulmonary cancer; in women, 65% of smoking-related deaths were also from this type of cancer.[16]

Environmental tobacco smoke (ETS), also known as second-hand smoke, has been classified by the International Agency for Research on Cancer as a Group 1 carcinogen, that is, there is convincing evidence that it produces cancer in humans.[17] A non-smoker's risk of contracting lung cancer has been shown to increase 20%–30% when exposed to ETS. ETS also produces an increase in all-cause death rates, as well as changes in the immune system.[17–20] Among the 41 countries that reported data on population exposure to ETS in 2002, Cuba ranked in the top five, and in the top 24 in a 2006 report on 82 countries.[21,22]

In 2001 in Villa Clara, 45%–49.9% of families were exposed to ETS. In Cuba, research indicates that persons with higher educational levels may tend to underestimate ETS risk, constituting failed responsibility for protection of their own and others' health. The extent of this problem in Cuba denotes a knowledge gap that could compromise future population health outcomes in the country.[23]

Cancer morbidity and mortality rates are indicators most frequently used in surveillance of cancer and lung cancer in Cuba. Variables considered risk factors (smoking, ETS exposure, work in certain industries, family history, etc.) may cluster in certain population groups located in certain places and at certain times. Thus, a population group's lifestyle and behaviors can determine exposure patterns, concentrating or excluding risk factors. Spatial aggregation or clustering can provide evidence of shared risk factors, and identify regions where strong patterns predominate of common ethnicity, eating habits, soil type, agricultural activities, climate, social organization, *inter alia*. [24]

For population health problems, characterizing their geographic and spatial differentiation reveals patterns that can provide important clues to causation and identification of systemic problems that may require intervention. Integrating such information into the public health surveillance system can offer important evidence for health service management and decisionmaking. Thus, consistent study of health problems' geographic distribution and assessment of inequalities in environmental and population living conditions is an additional tool for rational resource allocation and for achieving greater equity in health and all public policies.[25] Viewing geography and its spatial heterogeneity as context also helps link the conceptual framework of the health-illness process to how it plays out in reality.

The importance of identifying high-risk areas for intervention explains the current keen interest among epidemiologists in spatiotemporal distribution of diseases, and in particular, cluster analysis. A cluster is defined as the occurrence of a significantly greater number of cases than expected in a specific geographic area or specific time period, or both. Epidemiological demonstration of a cluster does not necessarily imply causal association between the possible risk factor and the disease. An etiologic study is required to confirm association when evidence indicates ill persons have been exposed to a common risk factor.[26]

Analyzing complex spatial and temporal patterns related to risk factors and disease has been used by several researchers in

the Americas, with encouraging results.[26–36] It has been infrequently applied to case fatality; and Cuba does not have a strong tradition of etiologic cancer research using spatial analysis.

We used spatial analysis to examine patterns of lung cancer morbidity and case fatality in Villa Clara Province to identify spatial and spatiotemporal clusters and demonstrate utility of this technique as an epidemiologic tool.

METHODS

A descriptive observational study was conducted in Villa Clara Province using a statistical scanning software program to explore spatial and spatiotemporal distributions of lung cancer.

The study reviewed records of patients residing in the province of Villa Clara aged >15 years who were diagnosed with lung cancer (387) during the period from January 1 to December 31, 2004; and among these, deaths occurring through 2009 (311).

Study variables Diagnosed cases, date of diagnosis, deaths, date of death, spatial units (in this case the municipalities in Villa Clara Province: Corralillo, Quemado de Güines, Sagua la Grande, Encrucijada, Camajuaní, Caibarién, Remedios, Placetas, Santa Clara, Cifuentes, Santo Domingo, Ranchuelo and Manicaragua), and Cartesian geocoding of each municipality.

Data source and analysis Data on lung cancer incidence in 2004 and deaths in these cases from 2004 through 2009 obtained from databases of the Provincial Health Statistics Bureau, Villa Clara. Incidence rates were calculated per 10,000 population using the number of lung cancer cases diagnosed in 2004 and the census population reported for that year. Case fatality was calculated as percentage of patients diagnosed in 2004 who died before the close of 2009.

To generate records required by the program on geographic coordinates, populations and cases, various databases were used:

- Information on lung cancer cases diagnosed in 2004, and the outcome of these through study termination on December 31, 2009, from Villa Clara's Provincial Health Statistics Bureau;
- Information on total population from the National Statistics Bureau; [7]
- Records on geocoding (Cartesian) of the province's municipalities from MapInfo v. 8.5.

To determine the existence of clusters by spatial analysis unit, Kulldorff's space-time scan statistic (SaTScan Program v. 7.01) was used.[37] The main advantages of this statistical tool include: it prevents selection bias; for each cluster a unique value of p is obtained, which is adjusted by the multiple tests derived from the large quantity of potential clusters evaluated; if the null hypotheses is rejected, the methodology indicates the approximate cluster site responsible.

Each type of analysis has a different purpose: spatial analysis disregards time and detects anomalous concentrations in space, while spatiotemporal analysis detects clusters coinciding in space and time. The space-time scan statistical method explained in detail by Kulldorff creates a type of circular window (space) or cylindrical window (space-time) with a circular base of varying radius, corresponding to a geographical area; the cylinder's height corresponds to time. These are centered

Original Research

around one of many possible centroids in the study, with each circle or cylinder representing a possible cluster. Each cylinder has a specific number of observed cases and the program calculates the number of expected cases under the null hypothesis for the total risk population.[38] We set α at 0.05 for rejecting the null hypothesis of 'no cluster.'

Relative risk (RR) estimates reported denote risk of incidence of cancer or mortality from cancer in the cluster, compared with risk in the rest of the province. Tables were created to display clusters with a statistically significant ($p < 0.05$) excess of observed over expected cases, under the hypothesis of a Poisson distribution and using the likelihood ratio for parameter estimation. Cartographic representations were made with MapInfo v. 8.5, importing statistical results into the geographic information system. Municipalities forming part of significant spatiotemporal clusters were colored and those in spatial clusters identified by a star.

RESULTS

Incidence Overall lung cancer incidence for the province was 4.8 per 10,000 population. Seven municipalities had higher rates: Caibarién (7.3), Camajuaní (6.3), Placetás (6.0), Sagua la Grande (5.9), Quemado de Güines (5.3), Encrucijada (5.1) and Cifuentes (5.0) (Table 1).


Spatial analysis Municipalities at highest risk are displayed in Figure 1. A maximum-likelihood spatial cluster was identified that included the municipalities of Encrucijada, Camajuaní, Cifuentes, Sagua and Caibarién, with 133 cases observed, compared to 106.5 expected ($p = 0.044$).

Spatiotemporal analysis A significant spatiotemporal cluster was detected that included the municipalities of Cifuentes, Encrucijada, Sagua la Grande and Santa Clara. During December 2004, there were 32 cases observed versus 14.5 expected ($p = 0.030$), that is, more than twice the risk observed in the rest of the province (Figure 1).

Table 1: Lung cancer incidence (2004) and case fatality (2004–2009), Villa Clara, Cuba

Municipality	Cases 2004 (No.)	Incidence (per 10,000 population)	Deaths 2004–2009 (No.)	Case fatality (%)
Corralillo	1	0.4	1	100.0
Quemado de Güines	12	5.3	9	75.0
Sagua la Grande	32	5.9	27	84.4
Encrucijada	17	5.1	14	82.4
Camajuaní	40	6.3	28	70.0
Caibarién	28	7.3	26	92.9
Remedios	17	3.7	13	76.5
Placetás	42	6.0	35	83.3
Santa Clara	107	4.5	78	72.9
Cifuentes	16	5.0	14	87.5
Santo Domingo	23	4.3	20	87.0
Ranchuelo	24	4.2	20	83.3
Manicaragua	28	3.9	26	92.9
Total	387	4.8	311	80.4

Figure 1: Spatial and spatiotemporal lung cancer incidence clusters in Villa Clara, Cuba (2004)



Analysis	Municipality	Time	Cases observed	Cases expected	RR	p Value
Spatial	Caibarién, Camajuaní, Cifuentes, Encrucijada, Sagua la Grande	—	133	106.5	1.37	0.044
Spatiotemporal	Cifuentes, Encrucijada, Sagua la Grande, Santa Clara	Dec 2004	32	14.5	2.31	0.030

Figure 2: Spatial and spatiotemporal lung cancer case fatality clusters in Villa Clara, Cuba (2004–2009)



Analysis	Municipality	Time	Deaths observed	Deaths expected	RR	p Value
Spatial	Caibarién, Camajuaní, Cifuentes, Encrucijada, Sagua la Grande	—	109	85.6	1.42	0.055
Spatiotemporal	Caibarién, Camajuaní, Cifuentes, Encrucijada, Sagua la Grande	Mar 1, 2004–Jun 30, 2005	87	19.0	5.96	0.001
	Manicaragua, Placetás, Santa Clara	Apr 1, 2004–Aug 31, 2005	104	34.5	4.03	0.001

Case fatality The provincial case fatality rate for 2004–2009 was 80.4% (Table 1). The municipalities with highest case fatality (not counting Corralillo, where only one case was diagnosed in 2004) were Caibarién and Manicaragua (92.9%), Cifuentes (87.5%) and Santo Domingo (87.0%).

Spatial analysis A single spatial cluster was found that included the municipalities of Encrucijada, Camajuaní, Cifuentes, Sagua la Grande and Caibarién, with a relative risk of 1.42 (Figure 2).

Spatiotemporal analysis The same municipalities as in the spatial cluster (Encrucijada, Camajuaní, Cifuentes, Sagua la Grande and Caibarién) formed a spatiotemporal cluster for the period of March 1, 2004–June 30, 2005, with 87 deaths versus 19 for an RR of 5.96. Another spatiotemporal cluster was identified in Manicaragua, Santa Clara and Placetas municipalities; the RR for the period April 1, 2004–August 31, 2005 was 4.03, with 104 deaths versus 34.5 expected (Figure 2).

DISCUSSION

One constraint in this investigation was the lack of similar studies in the national literature for comparison purposes. However, several international studies have applied this method to lung cancer. One was conducted by Fukuda, who identified the principal incidence clusters in Japan's metropolitan regions, related primarily to habits and socioeconomic conditions.[39] Another study conducted in Costa Rica identified high-incidence clusters of bronchopulmonary cancer in the country's northeast and Atlantic regions, implying a relationship with major risk factors such as smoking and environmental pollution.[26]

Incidence The lung cancer incidence clusters identified in this study indicate the need for further research on risk factors—mainly tobacco use, which constitutes the most significant etiological agent. A study conducted by Varona reported that some 23% of all deaths from cancer in Cuba in 2010 were due to lung cancer, of which 85% were attributed to smoking (92% in men and 75% in women).[40] An analytical cross-sectional epidemiological study in Matanzas Province, Cuba found that patients who smoked were four times more likely to die from lung cancer.[41]

Population aging is one of the most important contributors to rising numbers of cancer cases, and Villa Clara Province has the oldest population in the country (18.2% aged ≥ 60 years).[42] All of the province's municipalities are among the 40 oldest in the country (of a total of 169 municipalities).[7] Placetas, Cifuentes, Camajuaní and Encrucijada have the oldest populations in Villa Clara and are also where higher incidence and case fatality clusters were identified in this study.

Along with exposure to risk factors, genetic mutations predisposing to lung cancer may increase the probability of developing the disease. There is a hypotheses on varying susceptibility to carcinogens, based on an individual's ability to form genotoxic intermediates, detoxify these intermediates and repair the damage to DNA, all inheritable genetic features that would affect potential for suffering lung cancer and other cancers as well.[43]

Family aggregations can also be explained by the fact that individuals in the same family tend to have similar habits, such as smoking, drinking, diet and occupation, hence the value of studying clusters.[44]

The spatiotemporal incidence cluster in December 2004 could be due to year-end reporting procedures or other factors that warrant further examination in future studies. With a disease of such long latency, a biological explanation is implausible.

Case fatality Spatial analysis revealed close geographic overlap of incidence and case fatality (see Figures 1 and 2). However, case fatality patterns do not depend on incidence but are influenced more by stage at diagnosis and by treatment applied, both of which can affect patient survival.


Both spatiotemporal case fatality clusters were within two years of diagnosis, consistent with lung cancer's established poor prognosis, one of the worst for any cancer. It is asymptomatic over most of its evolution and, in most patients, by time of diagnosis, the disease is already well advanced and nonresponsive to most treatments.[40,45] Clearly screening and close examination of at-risk groups in primary health care are essential. This can overcome obstacles at various levels: individual (such as awareness), primary care (identification of at-risk individuals, streamlining of diagnosis and prompt referral to specialized care), and secondary care (prompt application of clinical protocol for diagnosed cases).[46] A study in Villa Clara Province found a direct association between patient survival and specific aspects of treatment: early tumor stage at diagnosis, morphologic and histologic diagnosis, and diagnostic methods used.[47] The finding that Placetas Municipality did not form part of any incidence clusters but did appear in a spatiotemporal case fatality cluster merits further study, as it could suggest health system factors at play, such as delays in diagnosis or errors in treatment protocols.

The importance of this type of study resides in its ability to raise questions and provide clues that may lead to causal explanations further down the road. Use of this tool in the surveillance system can help increase health equity and system efficiency by focusing medical services where they are most needed, and of course can help set priorities for research.

CONCLUSIONS

Application of the SaTScan analytical method revealed significant spatial and spatiotemporal lung cancer clusters, in both incidence and case fatality. While our findings are not explanatory, they do identify areas at higher risk and thus can support generation of hypotheses for research and eventual intervention.

Efforts are needed to determine factors contributing to the observed excess risk in areas of Villa Clara, to prevent the disease, initiate earlier detection and treat it more effectively, as well as to identify systemic factors that could account for differences in case fatality. Bringing about a major change in lung cancer burden will require improvements across the entire spectrum of prevention, early diagnosis and treatment.

It would also be valuable to extend application of this tool to epidemiological surveillance systems in other health jurisdictions in Cuba, to deepen epidemiological exploration of population health status, identify possible contextual determinants to facilitate better risk factor control, and seek solutions in health systems and services to problems identified through cluster detection. Such an investment would strengthen health and disease surveillance and provide a valuable addition to Cuba's universal health care system. 

REFERENCES

- World Health Organization. World Health Statistics [Internet]. 2008 [cited 2010 Feb 19]. Available from: <http://www.who.int/whosis/whostat/2008/en/>
- Pan American Health Organization. Plan de Acción Regional de Prevención y Control del Cáncer [Internet]. Washington DC: Pan American Health Organization; 2008 [cited 2012 May 23]. Available from: <http://www.paho.org/spanish/ad/dpc/nc/pcc-stakeholders-08.htm>. Spanish.
- Pan American Health Organization. Historia del cáncer, informes breves. *Bull Pan Am Health Organ*. 2007;121:577–8. Spanish.
- Cavalli F. Cáncer en el mundo en desarrollo: ¿podemos evitar el desastre? *Nat Clin Pract Oncol* [Internet]. 2006 [cited 2010 Dec 17];3(3). Available from: <http://scielo.sld.cu/scielo.php>. Spanish.
- Sansó F, Alonso P, Torres R. Mortalidad por cáncer en Cuba. *Rev Cubana Salud Pública*. 2010;36(1):78–94. Spanish.
- Anuario Estadístico de Salud 2011. Havana: Ministry of Public Health (CU); 2013. p. 41. Spanish.
- Anuario Estadístico Provincial. Villa Clara [Internet]. Havana: National Statistics Bureau (CU); 2009 [cited 2010 Jan 22]; [about 1 screen]. Available from: http://www.one.cu/publicaciones/provincias_masinf/villa%20Clara.htm. Spanish.
- Yassi A, Kjellström T, Kok T, Guidotti TL. Salud Ambiental Básica. Programa de las Naciones Unidas para el Medio Ambiente [Internet]. México DF: World Health Organization Regional Office for Latin America and the Caribbean; 2002 [cited 2009 Nov 10]. 551 p. Available from: <http://www.ambiente.gov.ar/infotecaea/descargas/yassi01.pdf>. Spanish.
- Pan American Health Organization. Evaluación de los efectos de la contaminación del aire en la salud de América Latina y el Caribe [Internet]. Washington DC: Pan American Health Organization; 2005 [cited 2009 Nov 10]. 42 p. Available from: http://www.bvssst.org.ve/documentos/portada/L238/efectos_contaminacion.pdf. Spanish.
- Biblioteca Virtual de Salud. Textos Completos [Internet]. Lima: Pan American Health Organization; 2013. Contaminación del aire en interiores. Una introducción para los profesionales de la salud; 1998 [cited 2009 Nov 10]; [about 2 screens]. Available from: <http://www.bvsde.paho.org/bvscie/fulltext/contint/contint.html>. Spanish.
- Barcelo C. Vivienda saludable por la construcción de la salud humana [Editorial]. *Rev Cubana Hig Epidemiol* [Internet]. 2007 [cited 2009 Nov 10];45(1). Available from: http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S156130032007000100001&lng=es. Spanish.
- Toh CK. The changing epidemiology of lung cancer. *Methods Mol Biol*. 2009;472:397–411.
- O'Reilly KMA, McLaughlin AM, Beckett WS, Sime PJ. Asbestos-related lung disease. *Am Fam Physician* [Internet]. 2007 [cited 2011 Apr 14];75(5):683–8. Available from: <http://www.aafp.org/atp/20070301/683.html>
- Kumar VRS, Cotran RS, Stanley LR. *Patología Humana*. 7th ed. Madrid: Elsevier Health Sciences; 2008. 500 p. Spanish.
- Varela S. El cáncer, epidemiología, etiología, diagnóstico y prevención. Madrid: Elsevier España; 2002. 165 p. Spanish.
- Varona P, Herrera D, García RG, Bonet M, Romero T, Venero SJ. Mortalidad atribuible al tabaquismo en Cuba. *Rev Cubana Salud Pública* [Internet]. 2009 Apr–Jun [cited 2011 Dec 13];35(2). Available from: http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S0864-34662009000200015&lng=es. Spanish.
- International Agency for Research on Cancer [Internet]. Geneva: World Health Organization; c2013. Agents Classified by the IARC Monographs. Volumes 1–107. 2013 [updated 2013 Apr; cited 2013 Jun 7]. 33 p. Available from: <http://monographs.iarc.fr/ENG/Classification/ClassificationGroupOrder.pdf>
- Hecht SS. Tobacco carcinogens their biomarkers and tobacco induced cancer. *Nat Rev Cancer* [Internet]. 2003 Oct [cited 2009 Nov 9];3(10):733–44. Available from: <http://toxicology.usu.edu/endnote/rr556SS2003.pdf>
- López MJ, Nebot M. La medición de la nicotina como marcador del humo ambiental de tabaco. *Gac Sanit* [Internet]. 2003 [cited 2009 Nov 10];17(3):15–22. Available from: <http://www.bvsde.paho.org/bvsacd/cd26/tabaco67.pdf>. Spanish.
- Mackay J, Eriksen M. The Tobacco Atlas [Internet]. Geneva: World Health Organization; 2002 Jun [cited 2009 Nov 9]. 54 p. Available from: <http://whqlibdoc.who.int/publications/2002/9241562099.pdf>
- Mackay J, Eriksen M, Shafey O. The Tobacco Atlas [Internet]. 2nd ed. Geneva: World Health Organization; 2006 [cited 2009 Nov 9]. Available from: http://whqlibdoc.who.int/publications/2006/ACS_Tobacco_Atlas_eng.pdf
- Varona P, García R, Molina E, Bonet M. Humo ambiental de tabaco en el hogar y percepción de riesgo en la población cubana. *Rev Cubana Hig Epidemiol*. 2010;48(3):291–303. Spanish.
- Alegret M. Propuesta metodológica para un uso más efectivo del análisis espacial en Ciencias de la Salud [thesis]. [Havana]: National School of Public Health (CU); 2006. 120 p. Spanish.
- Iñiguez L, Barcellos C. Geografía y salud en América Latina: evolución y tendencias. *Rev Cubana Salud Pública* [Internet]. 2003 [cited 2011 Jan 14];29(4):330–43. Available from: http://bvs.sld.cu/revistas/spu/vol29_4_031/spu07403.pdf. Spanish.
- Santamaría-Ulloa C. Análisis para la determinación de conglomerados espacio temporales en la incidencia de cáncer. Costa Rica, 1990–1997 [thesis] [Internet]. [San José]: Ciudad Universitaria Rodrigo Facio (CR); 2002 [cited 2010 Feb 8]. 84 p. Available from: http://ccp.ucr.ac.cr/bvp/pdf/tfgs2002/tfg_csantamaria.pdf. Spanish.
- Sánchez M. Denuncia de un posible conglomerado de casos en la Llanada de Rojas, Costa Rica. Informe oficial del Registro Nacional de Tumores. Dirección de Vigilancia de la Salud [Internet]. Costa Rica: Ministerio de Salud (CR); 2002 [cited 2011 Mar 14]. Available from: <http://ccp.ucr.ac.cr/revista>. Spanish.
- Santamaría-Ulloa C. El análisis espacial como herramienta para evaluar alarmas por cáncer. Costa Rica [Internet]. 2004 [cited 2011 Mar 20]. Available from: <http://ccp.ucr.ac.cr/revista>. Spanish.
- Joseph Sheehan T, DeChello LM, Kulldorff M, Gregorio DI, Gershman S, Mrosczyk M. The geographic distribution of breast cancer incidence in Massachusetts 1988 to 1997, adjusted for covariates. *Int J Health Geogr* [Internet]. 2004 Aug 3 [cited 2011 Mar 24];3(1):17. Available from: <http://www.ijhealthgeographics.com/content/3/1/17>
- Goovaerts P. Geostatistical analysis of disease data: estimation of cancer mortality risk from empirical frequencies using Poisson kriging. *Int J Health Geogr* [Internet]. 2005 Dec 14 [cited 2011 Mar 24];4:31. Available from: <http://www.ij-healthgeographics.com/content/4/1/31>
- Jacquez GM, Greiling DA. Local clustering in breast, lung and colorectal cancer in Long Island, New York. *Int J Health Geo* [Internet]. 2003 Feb 17 [cited 2011 Mar 31];2(1):3. Available from: <http://www.ij-healthgeographics.com/content/2/1/3>
- Collado A. Distribución espacial de la fecundidad adolescente en la gran área Metropolitana de Costa Rica [Internet]. San José (CR): University of Costa Rica; 2000 [cited 2011 Apr 4]. 39 p. Available from: <http://www.ccp.ucr.ac.cr/noticias/simposio/pdf/collado.pdf>. Spanish.
- Alegret M, Herrera M, Grau R. Las técnicas de estadística espacial en la investigación salubrista. Caso síndrome de Down. *Rev Cubana Salud Pública* [Internet]. 2008 Dec [cited 2011 Feb 12];34(4). Available from: http://www.bvs.sld.cu/revistas/spu/vol34_4_08/spu03408.htm. Spanish.
- Kulldorff M. Prospective time-periodic geographical disease surveillance using a scan statistic. *J R Statist Soc* [Internet]. 2001 [cited 2010 May 14];64(P1):61–72. Available from: <http://satscan.us/papers/k-jrssa2001.pdf>
- Neill DB, Cooper GF. A multivariate Bayesian scan statistic for early event detection and characterization. *Mach Learn* [Internet]. 2009 Nov 13 [cited 2010 May 14];79(3):261–82. Available from: <http://www.cs.cmu.edu/~neill/papers/MBSS.pdf>
- Kulldorff M, Athas WF, Feurer EJ, Miller BA, Key CR. Evaluating cluster alarms: A space-time scan statistic and brain cancer in Los Alamos. *Am J Public Health* [Internet]. 1998 Sep [cited 2010 May 14];88(9):1377–80. Available from: http://www.researchgate.net/publication/13548701_Evaluating_cluster_alarms_a_space-time_scan_statistic_and_brain_cancer_in_Los_Alamos_New_Mexico
- Kulldorff M. Information Management Services, Inc. SaTscanTM v7.0: Software for the spatial and space-time scan statistics [Internet]. Boston: SaTscan; 2005 [cited 2010 Mar 14]. Available from: <http://www.satscan.org/>
- Kulldorff M. A spatial scan statistic. *Commun Statistics: Theory and Methods* [Internet]. 1997 [cited 2011 Mar 14];26(6):1481–96. Available from: <http://www.tandfonline.com/doi/abs/10.1080/03610929708831995#preview>
- Fukuda Y, Umezaki M, Nakamura K, Takano T. Variations in societal characteristics of spatial disease clusters: examples of colon, lung and breast cancer in Japan. *Int J Health Geogr* [Internet]. 2005 Jun 14 [cited 2011 Mar 8];4:16. Available from: <http://www.ij-healthgeographics.com/content/4/1/16>
- Varona P, Torres P, Elejalde AR, Hernández EA, Neninger E. Modelo para la prevención y manejo del cáncer de pulmón en Cuba, 2010. *Rev Cubana Hig Epidemiol* [Internet]. 2012 Jan–Apr [cited 2013 Apr 3];50(1):37–47. Available from: http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S1561-30032012000100006&lng=es. Spanish.
- Achiong F, Morales JM, Dueñas A, Acebo F, Bermúdez C, Garrote I. Prevalencia y riesgo atribuible al tabaquismo. *Rev Cubana Hig Epidemiol* [Internet]. 2006 [cited 2011 Oct 5];44(1). Available from: http://www.bvs.sld.cu/revistas/hie/vol44_1_06/hie01106.htm. Spanish.
- Rodríguez A, Álvarez L. Repercusiones del envejecimiento de la población cubana en el sector salud. *Rev Cubana Salud Pública* [Internet]. 2006 Apr–Jun [cited 2013 Jan

- 21];32(2):178–82. Available from: http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S0864-34662006000200013&lng=es. Spanish.
43. Sridhar SS, Seymour L, Shepherd FA. Inhibitors of epidermal-growth-factor receptors: a review of clinical research with a focus on non-small-cell lung cancer. *Lancet Oncol*. 2003 Jul;4(7):397–406.
44. Reyes K, Marcheco B, Crombet T. Agrupación familiar para el cáncer en individuos afectados por cáncer de pulmón. *Rev Cubana Genet Comunit*. 2010;4(1):11–8. Spanish.
45. Huet J, Cebollero P, Cascante JA, Andrade I, Pascal I, Boldú J, et al. Evaluación de la utilización de una consulta de diagnóstico rápido de cáncer de pulmón. *Tiempos de demora diagnóstica y terapéutica*. *Rev Arch Bronconeumol [Internet]*. 2012 [cited 2013 Apr 3];48:267–73. Available from: <http://www.sciencedirect.com/science/article/pii/S030028961200107X>. Spanish.
46. Monteagudo A, Romero RE, Salazar CE. Cáncer de pulmón, propuesta ética para agilizar su diagnóstico. *Rev Hum Med [Internet]*. 2007 May–Aug [cited 2011 Mar 14];7(2). Available from: http://scielo.sld.cu/scielo.php?pid=S1727-81202007000200004&script=sci_arttext. Spanish.
47. Hernández D, Alegret M. Supervivencia en el cáncer pulmonar: una necesidad de los servicios de salud en Villa Clara. *Rev Medicentro Electrón [Internet]*. 2012 Jul–Sep [cited 2013 Apr 3];16(3) [about 3 screens]. Available from: <http://medicentro.troajs.vcl.sld.cu/index.php/medicentro/article/viewFile/1276/1200>. Spanish.

THE AUTHORS

Norma E. Batista Hernández (Corresponding author: normabh@ucm.vcl.sld.cu), physician specializing in biostatistics and family medicine, biomedical research unit, Medical University of Villa Clara, Santa Clara, Cuba.

Oscar A. Antón Fleites, family physician, Marta Abreu University Teaching Polyclinic, Santa Clara, Cuba.

Submitted: January 5, 2012

Approved for publication: July 12, 2013

Disclosures: None

PSICO Habana 2014

Havana Psychiatric Hospital's 6th International Conference

Havana's International Convention Center
March 17–21, 2014

Themes

- Clinical psychiatry and psychology
- Psychiatry research and classification
- Addictions
- Pharmacotherapy for mental disorders
- Forensic psychiatry and psychology
- Psychiatric rehabilitation
- Nursing
- Psychoballet

Sponsors

- Ministry of Public Health (CU)
- Council of Scientific Societies in Health (CU)
- Buenos Aires Mental Health Cooperative

Conference language: Spanish * Contact: psico2014@infomed.sld.cu * www.psicohabana.com