

Epidemiology of snakebites in Colombia (2008-2016)

Epidemiología de las mordeduras por serpientes en Colombia (2008-2016)

Leonardo J. León-Núñez, Gabriel Camero-Ramos and José M. Gutiérrez

Received 5th May 2020 / Send for modification 5th June 2 2020 / Accepted 5th June 2020

ABSTRACT

Objective To describe the main epidemiological features of snakebites in Colombia during the period 2008 to 2016.

Methods A retrospective (quantitative) descriptive analytical empirical study was carried out, based on the official databases of the Public Health Surveillance in the Integral Information System of the Social Protection (SISPRO) and the Surveillance System in Public Health (Sivigila) of the reported cases of snakebites in Colombia for that period.

Results In total, 37 066 cases were reported, with annual incidences ranging from 7.0 (2008) to 9.7 (2011 and 2012) cases per 100,000 population. Mortality rates ranged from 0.059 (2013) to 0.091 (2011) deaths per 100 000 population, with case fatality rates ranging from 0.6% (2013) to 1.0% (2010). Indigenous and Afro-Colombian populations were highly affected, and highest incidences occurred in males, and in people living in rural areas. The average age of affected people is 31.7 years (95% CI 28.3 34.5). The regions with higher incidence are Amazonia and Orinoquia. Species of the genus *Bothrops* are responsible for the highest number of bites (64.5%), owing to their wide distribution in Colombia. Regarding clinical manifestations, pain and edema were observed in 86.9% and 78.8% of patients, respectively. Cellulitis and abscesses were the most frequent local complications of these envenomings.

Conclusions Results underscore the relevance of snakebite envenoming in Colombia, and provide information for improving the public health attention to these envenoming.

Key Words: Snakebite envenoming; Colombia; epidemiology; incidence; mortality; *bothrops* (source: MeSH, NLM).

RESUMEN

Objetivo Describir las principales características epidemiológicas de las mordeduras por serpientes en Colombia durante el período 2008-2016.

Métodos Se realizó un estudio retrospectivo (cuantitativo) analítico empírico sobre los reportes de accidentes por mordedura de serpientes en Colombia para el período 2008 a 2016, a partir de las bases de datos oficiales del Sistema Integral de Información de la Protección Social (SISPRO) y del Sistema Nacional de Vigilancia en Salud Pública (Sivigila).

Resultados Se reportaron 37 066 casos de mordeduras por serpientes en ese período. Las incidencias anuales oscilaron entre 7,0 (2008) y 9,7 (2011 y 2012) casos por 100 000 habitantes. Las tasas de mortalidad oscilaron entre 0,059 (2013) y 0,091 (2011) muertes por 100 000 habitantes, con tasas de letalidad entre 0,6% (2013) y 1,0% (2010). Los grupos más afectados fueron las poblaciones indígenas y afrocolombianas, y las personas residentes en zonas rurales. La edad promedio de las personas afectadas fue 31,7 años (IC 95% 28,3 34,5). Las regiones con mayor incidencia fueron la Amazonía y la Orinoquia. Especies del género *Bothrops* fueron responsables del mayor número de casos (64,5%), debido a su amplia distribución en el territorio colombiano. Con relación a las manifestaciones clínicas, se describió dolor y edema en 86,9% y 78,8% de los pacientes,

LL: MD. Enf. Esp. Epidemiología.
M. Sc. Estudios de Población. Cruz Roja,
Seccional Cundinamarca y Bogotá. Bogotá,
Colombia.

lj.leon62@uniandes.edu.co

GC: MD. Esp. Epidemiología.

Esp. Salud Familiar. Esp. Estadística.

M. Sc. Educación Salud. FETP Training Programs
in Epidemiology and Public Health Interventions
Network TEPHINET Cruz Roja, Seccional
Cundinamarca y Bogotá. Bogotá, Colombia.

gabriel.camero@gmail.com

JG: Lic. Microbiología y Química Clínica.

Ph. D. Ciencias Fisiológicas; Instituto Clodomiro
Picado. Facultad de Microbiología,

Universidad de Costa Rica. San José, Costa Rica.

jose.gutierrez@ucr.ac.cr

respectivamente. Las complicaciones locales más frecuentes fueron las celulitis y los abscesos.

Conclusiones Los resultados muestran la importancia de las mordeduras por serpientes en Colombia y suministran información para mejorar la atención de la salud pública ante estos envenenamientos.

Palabras Clave: Mordedura de serpiente; Colombia; epidemiología; incidencia; mortalidad; bothrops (*fuentes: DeCS, BIREME*).

Snakebite envenoming is a public health problem of high impact on a global basis, having its heaviest burden in sub-Saharan Africa, Asia and Latin America (1) and mostly affecting rural impoverished populations in the rural tropics (2). In 2017, the World Health Organization (WHO) included snakebite envenoming as a category A disease in its list of neglected tropical diseases. Moreover, a resolution on snakebite envenomings was adopted by the World Health Assembly in 2018, and the WHO launched its global strategy for prevention and control of this disease in 2019 (3). These initiatives aim at reducing the number of deaths and disabilities due to snakebite envenomings by 50% by the year 2030 (3). A key issue in the fulfillment of this global strategy is the acquisition of robust epidemiological information on snakebites in countries with high incidence.

It has been estimated that the total annual number of snakebite envenomings in the world ranges from 1.8 to 2.7 million cases, resulting in 81,000 to 138,000 deaths and in more than 400,000 people left with permanent disabilities (1). In the Americas, an overall incidence and mortality rate of 6.2 and 0.04 per 100,000 population per year, respectively, were described, with a case fatality rate of 0.64% (4). Snake species of the families *Viperidae* and *Elapidae* are responsible for the most severe cases of envenoming in the world, with fewer cases inflicted by species of the family *Lamprophiidae* and the non-front fanged *Colubroid* families (1). In Colombia, the report of snakebite envenoming as a single entity started in the year 2008. Previous reports in this country indicate incidences of 6 to 8.5 cases per 100,000 population per year, with a case fatality rate ranging from 1 to 3% (5). More recently, and based on various sources of epidemiological information, Chippaux (4) reported for Colombia, in the period 2009 to 2014, incidence and mortality rate of 8.69 and 0.073 per 100,000 population per year, respectively, with a case fatality rate of 0.84%. The Instituto Nacional de Salud (INS) of Colombia reported 4978 cases of snakebites in the year 2017 (6). Other epidemiological studies have described a relationship between snakebite incidence and precipitation in almost 50% of the municipalities in this country (7). Previous reports have focused on particular regions in Colombia, such as Antioquia and Chocó (8), Nariño (9), and the Coffee Triangle Region (10).

Colombia has an abundant herpetofauna of venomous snakes, the most important of which are classified in the

families *Viperidae* and *Elapidae*. The species of maximum medical impact are the viperids *Bothrops asper*, *B. atrox*, *B. bilineatus* and *Crotalus durissus* (5). In addition, other *Bothrops* species, and species classified in the genera *Bothriechis*, *Bothrocophias*, *Lachesis*, and *Porthidium* (family *Viperidae*), and coral snakes of the genus *Micrurus* (family *Elapidae*) also cause human envenomings (5).

It is necessary to have national-wide information on the main epidemiological features of snakebites in Colombia in order to have an integrated view of the characteristics of this disease in the country. The present investigation was undertaken to analyze the epidemiology and some clinical features of this neglected tropical disease in Colombia for the period 2008-2016. This information provides a base of knowledge for the design and implementation of public health policies in this country aimed at developing cost-effective interventions to confront this disease.

METHODO

A transversal, retrospective descriptive study was done. The data bases used were the official data of Public Health Surveillance in the Integral Information System of the Social Protection (SISPRO) and the Surveillance System in Public Health (Sivigila) of Colombia, for the period 2008-2016. For the estimation of indicators based on population, the municipal area projection 1985-2020, from the National Department of Statistics (DANE), was used. Annual snakebite incidences and mortality rates were calculated by using the DANE data on population of each year, and the incidences per department were also estimated. The identification of the types of snakes responsible for the accidents, as well as the types of localities where bites occurred, the type of health affiliation system through which accidents were treated, and the main clinical manifestations of envenomings, were also collected from these data bases. The estimation of descriptive information, the trend curves and the correlation between various parameters were carried out with the program Excel. Parametric (Student's *t* test, correlation of Pearson, and Chi squared) and non-parametric (Mann-Whitney) tests were used in the analyses depending on the type of variable, case and groups analyzed. Some results were expressed with the 95% confidence intervals (95% CI). Values of $p < 0.05$ were considered significant.

RESULTS

In the interval 2008 to 2016, a total of 37,066 cases of snakebites were reported in Colombia, with a range of 38 to 120 cases per epidemiological week. The total number of cases per year ranged from 3126 (year 2008) to 4704 (year 2016), and the number of fatalities ranged from 28 (year 2008) to 42 (year 2011). In turn, case fatality rates ranged from 0.6% (year 2013) to 1.0% (year 2010) (Table 1). Annual incidences ranged from 7.0 (2008) to 9.7 (2011 and 2012) cases per 100,000 population. Mortality rates ranged from 0.059 (2013) to 0.091 (2011) per 100,000


population (Table 1). There was a trend for increase in the number of bites along this period, with an R2 value of 0.82, with peaks of incidence for the years 2011, 2012 and 2016 (Figure 1). The highest incidences occurred in males, also with an incremental trend (R2 0.7), with male/female ratios ranging from 2.4 to 2.7 along this period (Figure 2). The mean age of people suffering snakebites in Colombia in this period was 31.7 years (95% CI: 28.3-34.5), with a median age of 28.9 years. The majority of cases (70.3%) correspond to males, and 34.7% of the cases were males younger than 30 years old (Figure 2).

Table 1. Incidence and mortality due to snakebite envenomings in Colombia (2008-2016)

Year	2008	2009	2010	2011	2012	2013	2014	2015	2016
Cases of snakebites	3126	3405	3945	4455	4526	4400	4232	4273	4704
Deaths	28	31	38	42	38	28	34	31	34
Incidence per 100,000 population	7.0	7.6	8.7	9.7	9.7	9.3	8.9	8.8	9.6
Mortality rate per 100,000 population	0.063	0.069	0.083	0.091	0.082	0.059	0.071	0.064	0.070
Case fatality rate (%)	0.9	0.9	1.0	0.9	0.8	0.6	0.8	0.7	0.7


Source: Data from SISPRO and Sivigila, Colombia, 2008 -2016.

Figure 1. Number of snakebite cases in Colombia (A) and incidences per year (B) for the period 2008-2016



Source: Public Health Surveillance in the Integral Information System of the Social Protection (SISPRO) and the Surveillance System in Public Health (Sivigila) of Colombia.

Figure 2. Distribution of snakebites occurring in Colombia during the period 2008-2016 according to age, for males and females and for urban and rural locations




Source: Public Health Surveillance in the Integral Information System of the Social Protection (SISPRO) and the Surveillance System in Public Health (Sivigila) of Colombia.

The highest number of cases and incidences were in rural regions (Figure 2), with a peak of 31.9 per 100,000 population for the year 2016 and with an incremental trend over the period ($R^2=0.79$), whereas the overall incidence in the period for urban areas was 2.8 per 100,000 population per year. Snakebites occurred throughout the country, with the exception of San Andrés, a Caribbean island. The geographical regions of Amazonia and Orinoquia presented the highest incidences, particularly the departments of Amazonas, Vaupés, Guaviare, Vichada and Arauca (Figure 3). Cases occurred throughout the

year, without drastic differences in the number of snakebites along the different months (not shown). When cases were classified according to the self-declared ethnic groups, Afro-Colombian and indigenous people showed the highest percentages among self-declared groups, corresponding to 9.9% and 9.7% of the cases, respectively, whereas most affected people (79.6%) were classified as 'others' in terms of ethnic group, and percentages of less than 1% corresponded to roma, raizel and palenquero groups.


Figure 3. Mean annual incidence of snakebites in Colombia per 100,000 population per departments for the period 2008-2016



The departments correspond to: (1) Vaupés; (2) Amazonas; (3) Guaviare; (4) Arauca; (5) Vichada; (6) Casanare; (7) Putumayo; (8) Chocó; (9) Caqueta; (10) Guainía; (11) Meta; (12) César; (13) Bolívar; (14) Norte de Santander; (15) Sucre; (16) Magdalena; (17) Córdoba; (18) Huila; (19) La Guajira; (20) Antioquia; (21) Cauca; (22) Santander; (23) Atlántico; (24) Nariño; (25) Caldas; (26) Tolima; (27) Boyacá; (28) Risaralda; (29) Quindío; (30) Valle del Cauca; (31) Cundinamarca; (32) San Andrés. Elaboration of map: Dany Villalobos.

In Colombia, health services are provided according to the Social Security General System in Health (*Sistema General de Seguridad Social en Salud, SGSSS*). People are ascribed to various affiliation systems. Those who pay to receive health services belong to a contributive system, whereas those who are unable to cover the costs of the health service are subsidized by the public health system. In addition, groups classified as ‘special’ and ‘exception’ mainly belong to the armed forces and the education sector. According to this classification, the majority of people affected by snakebites in this period belong to the subsidized regime (70.4%), followed by non-insurance (15.5), contributive (11.0%), special (2.2%), exception (0.7%), and undetermined (0.2%).


Figure 4. Percentage distribution of cases of snakebites in Colombia, on the basis of the genus of the snake causing the accident, for the period 2008-2016



NI: Offending snake was not identified. Others: Snakes of other genera, such as *Bothriechis* and *Porthidium*. Source: Public Health Surveillance in the Integral Information System of the Social Protection (SISPRO) and the Surveillance System in Public Health (Sivigila) of Colombia.

In Colombia, the severity of snakebite envenomings is classified as no envenoming, mild, moderate and severe envenoming, according to the local and systemic clinical manifestations of envenoming and the alterations in clinical laboratory parameters, depending on the genus of the


Figure 6. Local clinical manifestations in snakebite envenomings occurring in Colombia in the period 2008-2016



Source: Public Health Surveillance in the Integral Information System of the Social Protection (SISPRO) and the Surveillance System in Public Health (Sivigila) of Colombia.

Regarding the identity of the offending snake, according to the data collected in the information systems analyzed, snakes of the genus *Bothrops* inflicted 64.5% of the cases (Figure 4). In terms of the activity people were doing at the time of the bite, the majority of the cases occurred while carrying out agricultural work, followed by domestic work (in rural settings), and walking in rural trails (Figure 5). The early management of snakebite cases included, in 30.8 % of the cases, a variety of interventions outside health facilities, such as praying, oral administration or topical application of plant extracts, suction with the mouth at the site of the bite, and attendance by local healers. In 73.5% of the cases antivenom was administered at health centers.


Figure 5. Percentage distribution of cases of snakebites in Colombia, on the basis of the activity being carried out at the time of the bite, for the period 2008-2016



Source: Public Health Surveillance in the Integral Information System of the Social Protection (SISPRO) and the Surveillance System in Public Health (SIVIGILA) of Colombia.

offending snake (11). On this basis, 61.5% of the cases were classified as mild envenomings; among them, 67.6% received antivenom. On the other hand, 32.2 % of cases were classified as moderate, and 6.3% as severe. Of these, 85.3% and 87.7%, respectively, received antivenom as part

Figure 7. Systemic clinical manifestations in snakebite envenomings occurring in Colombia in the period 2008-2016



Source: Public Health Surveillance in the Integral Information System of the Social Protection (SISPRO) and the Surveillance System in Public Health (Sivigila) of Colombia.

of the treatment. The local and systemic clinical manifestations of envenomings are depicted in Figures 6 and 7.

The most common local complications of envenomings were associated with infection (19.5% of cases with cellulitis and 2.9% with abscesses). In terms of systemic complications, anemia was described in 1% of cases, and acute kidney injury in 0.8%. Surgical interventions were carried out in 1029 cases, including surgical cleansing procedures (39.75%) and drainage of abscesses (33.72%).

DISCUSSION

This study presents an overview of the main epidemiological features of snakebites in Colombia from 2008 to 2016. The data examined reveal a general trend of an increment in reported snakebites in Colombia over the period 2008-2016. This could reflect an actual increase in the number of bites in the country, but more likely is the consequence of the improvements in the notification of cases. The *Instituto Nacional de Salud* (INS) and other stakeholders developed a series of workshops in various regions of Colombia from 2015 to 2018, which is likely to have impacted in the report of the cases. Other possible explanations can be considered as well. For instance, changes in the incidence of snakebites may be related to climate parameters, as described in Costa Rica (12), demographic changes, in-country migrations, and changes in the use of the land. In several municipalities in Colombia, a relationship between snakebite incidence and precipitation has been demonstrated (7). It is necessary to carry out more detailed studies to discern which factors are involved behind this increment in the reported number of snakebite cases in Colombia. Despite the described improvement in the collection of data on snakebites, it is likely that under-registration still occurs, especially in regions where snakebites are not attended in health facilities, and this is a limitation of our study. There is a need to identify such regions in order to improve the official records of this neglected tropical disease.

Incidence of snakebites in the period of study in Colombia ranged from 7.0 and 9.3 per 100,000 population per year. These values agree with the value of 8.69 per 100,000 population described by Chippaux (4) for the period 2009-2014. A similar incidence has been described in other countries of Latin America, although some countries in the region show higher incidences, such as Panama, Guyana and French Guyana (4). In terms of mortality rate in the period studied, Colombia had rates that ranged between 0.059 (2013) and 0.091 (2011) deaths per 100,000 population per year. Chippaux (4) described a mean mortality rate for the period 2009-2014 of 0.073 deaths per 100,000 population per year.

Mortality rates in Latin America ranged from 0.012 (Argentina) to 0.631 (French Guyana) per 100,000 population per year (4). When case fatality rates are analyzed, our study showed a range of 0.7% to 1.0%, in agreement with the value of 0.84% described for the period 2009-2014 (4). In Latin America, for this period, case fatality rates ranged from 0.42% (Brazil) to 4.44% (Bolivia) (4). In the Brazilian Amazonia, higher mortality has been described among older people and people from indigenous communities with difficulties for having access to health services and antivenoms (13). It is relevant to investigate whether these factors are also associated with higher mortality in Colombia.

The regions with highest incidences of snakebites in Colombia are the Amazonia and Orinoquia, with highest incidences in the departments of Vaupés, Amazonas, Guaviare, Arauca, Casanare and Vichada. In agreement, the Amazon region in Brazil presents a high incidence of snakebites (14). These departments in Colombia have eco-epidemiological characteristics associated with high incidences of snakebites, i.e. climatic conditions that favor a rich snake fauna, and human activities characterized by agricultural work. A relationship has been found in Colombia between incidence of snakebites and precipitation in about 50% of the municipalities, but not in others (7). Likewise, in the Brazilian Amazonia, snakebites are more frequent in regions of high precipitation and humidity (15). This combination of ecological features and human activities increases the likelihood of encounters between venomous snakes, especially of the genus *Bothrops*, and humans, as described for the rest of Latin America (16). The identification of the departments and regions of highest incidence of snakebites in Colombia, and the analysis of the access to health posts in these settings (see for example (17)), is relevant for the design of public health interventions tailored to the characteristics and needs of those regions, hence making interventions more effective. This has been stressed in the WHO strategy for prevention and control of snakebite envenomings (3).

Snakebites are more frequent in rural areas than in urban settings. The activity associated with the highest number of snakebite cases was agricultural work, as previously described in Colombia (5) and in many regions of the world (1). In this context males are more affected than females, particularly young males in their productive age. Therefore, the study of changes in the use of land in rural areas is relevant for identifying changing patterns in snakebite incidence. Examples of such changes are deforestation for expanding agricultural land or creating new urban settings, as well as for development of mining activities. Similar findings of high incidence of snakebites in agricultural settings have been described for Ecuador

(18), Costa Rica (19), Bolivia (20), and Brazil (21), among other countries in the region. Despite this trend, it is relevant to identify other circumstances in which snakebites occur. Our work found domestic activities or walking through rural trails as common scenarios of snakebites. *Bothrops asper* and *B. atrox*, two species responsible for the highest number of cases in Colombia (5,11, this work), are ecologically-versatile species (22) which can be found nearby human dwellings, hence explaining the occurrence of bites near or inside houses.

There are various ways of affiliation to the Health Social Security System in Colombia. The highest numbers of snakebite cases correspond to people of the low income of the most vulnerable sectors of the population, and these are mostly attended through the subsidized system of public health. It is relevant to identify human groups which are particularly prone to suffer snakebites. Such is the case of indigenous communities in Latin America, which have a high incidence of snakebites and often lack appropriate health services (23,24). The relationship between health issues and ethnicity has been highlighted by the Pan American Health Organization (25). Interventions tailored to the cultural contexts of these groups should be designed and implemented in Colombia and in other countries in the region.

The majority of cases in which the offending snake was identified corresponded to *Bothrops* sp bites, most likely *B. asper* or *B. atrox*. This agrees with the general trend described in Colombia (5). Fewer cases were inflicted by rattlesnakes (*Crotalus durissus*) and the bushmaster (*Lachesis muta* and *L. acrochorda*), and by coral snakes of the genus *Micrurus*. Envenomings by *C. durissus* differ from those of other viperids for being predominantly neurotoxic and myotoxic (5). However, it is more difficult to differentiate between cases inflicted by *Bothrops* sp and *Lachesis* sp, since they have similar clinical manifestations. Nevertheless, bites by *Lachesis* are less frequent owing to its predominant distribution in primary forest, whereas *B. asper* and *B. atrox* adapt well to agricultural settings and other altered settings (22).

The data collected in the information systems used in this study allow for the analysis of the main clinical features of envenomings. In terms of the severity of the clinical manifestations, cases were classified as mild, moderate or severe in 61.5%, 32.2% and 6.3%, respectively. A more detailed characterization of the severity of the cases has been proposed by Otero-Patiño (11) who graded severity in terms of local and systemic manifestations. According to the clinical manifestations described, the main local effects were pain and edema, in agreement with previous studies describing clinical features in envenomings by *Bothrops* sp (5,11,19,26,27). In terms of systemic effects,

nausea and vomiting were often described, and constitute typical manifestations of viperid envenomings (27). In general, the percentage of patients with systemic manifestations in this study was lower than in previous reports (5,11). This discrepancy reveals limitations in the report of clinical manifestations in the notification systems used in our study and underscores the relevance of using hospital files and prospective clinical studies for detailed analyses of clinical features of envenomings. The most common complications described were associated with local infections, i.e. cellulitis and abscess formation, and necrosis, in agreement with previous works (11,28). Local necrosis of soft tissues may end up in permanent sequelae, with a high impact in the quality of life of people affected (28). A low percentage of cases were associated with respiratory failure, probably caused by bites of coral snakes (*Micrurus* sp) or rattlesnakes (*Crotalus durissus*).

In terms of antivenom treatment, among patients whose envenomings were graded as mild, moderate or severe, the percentages that received antivenom therapy were 68%, 85% and 88%, respectively. The fact that more than 10% of patients having moderate or severe envenomings did not receive antivenom urges attention, as to identify the causes of this observation. Antivenoms from various public and private laboratories are distributed in Colombia, and the recommended dosage varies depending on the neutralizing potency of each antivenom (5). The issue of antivenom availability and accessibility in the various regions of Colombia deserves more detailed investigations.

In conclusion, our study summarizes the main epidemiological features of snakebite envenoming in Colombia for the period 2008-2016. Snakebites constitute the main type of accident caused by venomous animals in this country (29). The information presented in our work allows for the identification of trends in the occurrence of snakebites and for highlighting the regions and human groups at highest risk of suffering snakebites. This, in turn, provides a valuable base of knowledge for the design and implementation of public health interventions, including prevention, early management of cases and appropriate attention in health facilities. More detailed studies are required to further identify regions and groups at high risk, and ways to improve the provision of health services in these and other settings ♣

Acknowledgments: The collaboration of Dany Villalobos in the preparation of the map of incidences per department is greatly acknowledged.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflicts of interest: None.

REFERENCES

- Gutiérrez JM, Calvete JJ, Habib AG, Harrison RA, Williams DJ, Warrell DA. Snakebite envenoming. *Nat Rev Dis Primers*. 2017; 3:17079. DOI:10.1038/nrdp.2017.79.
- Harrison RA, Hargreaves A, Wagstaff SC, Faragher B, Laloo DG. Snake envenoming: a disease of poverty. *PLoS Negl Trop Dis*. 2009; 3:e569. DOI:10.1371/journal.pntd.0000569.
- World Health Organization. Snakebite Envenoming. A Strategy for Prevention and Control. Geneva: World Health Organization; 2019.
- Chippaux JP. Incidence and mortality due to snakebite in the Americas. *PLoS Negl Trop Dis*. 2017; 11:0005662. DOI:10.1371/journal.pntd.0005662.
- Otero-Patiño R. Snake bites in Colombia. In: Gopalakrishnakone P. *Clinical Toxinology*. Dordrecht: Springer Science; 2014. DOI:10.1007/978-94-007-6288-6_41-2.
- Instituto Nacional de Salud. Informe evento de accidente ofídico, período epidemiológico I-XII, Colombia, 2018. Bogotá: Instituto Nacional de Salud; 2018.
- Angarita-Gerlein D, Bravo-Vega CA, Cruz C, Forero-Muñoz NR, Navas-Zuloaga MG, Umaña-Caro JD. Snakebite dynamics in Colombia: Effects of precipitation seasonality on incidence. *Int Res Exp Students*. 2017:1-4. Available from: <https://bit.ly/3i2pCcp>.
- Otero R, Tobón GS, Gómez L, Osorio RG, Valderrama R, Hoyos D, et al. Accidente ofídico en Antioquia y Chocó. Aspectos clínicos y epidemiológicos (marzo de 1989-febrero de 1990). *Acta Médica Colombiana*. 1992; 17:229-49. Available from: <https://bit.ly/3jVafTw>.
- Sevilla-Sánchez MJ, Mora-Obando D, Calderón JJ, Guerrero-Vargas JA, Ayerbe-González S. Accidente ofídico en el departamento de Nariño, Colombia: Análisis retrospectivo, 2008-2017. *Biomédica*. 2019; 39:715-36. Available from: <https://bit.ly/3jScCqj>.
- Patiño-Barbosa AM, Herrera-Giraldo AC, Lozada-Riascos CO, Paniz-Mondolfi AE, Suárez JA, Rodríguez-Morales AJ. Snakebites mapping in municipalities of the Coffee Triangle region in Colombia using Geographic Information Systems (GIS). *Rev Panam Enf Inf*. 2019; 21:14-20. Available from: <https://bit.ly/3h6YHuS>.
- Otero-Patiño R. Epidemiological, clinical and therapeutic aspects of *Bothrops asper* bites. *Toxicon*. 2009; 54:998-1011. DOI:10.1016/j.toxicon.2009.07.001.
- Chaves LF, Chuang T, Sasa M, Gutiérrez JM. Snakebites are associated with poverty, weather fluctuations, and El Niño. *Sci Adv*. 2015; 1:e1500249. DOI:10.1126/sciadv.1500249.
- Da Silva Souza A, Sachett JAG, Alcántara JA, Freire M, Alecrim MDGC, Lacerda M, et al. Snakebites as cause of death in the Western Brazilian Amazon. Why and who dies? Deaths from snakebites in the Amazon. *Toxicon*. 2018; 145:15-24. DOI:10.1016/j.toxicon.2018.02.041.
- Feitosa EL, Sampaio VS, Salinas JL, Queiroz AM, da Silva IM, Gomes AA, et al. Older age and time to medical assistance are associated with severity and mortality of snakebites in the Brazilian Amazon: A case-control study. *PLoS One*. 2015; 10:e0132237. DOI:10.1371/journal.pone.0132237.
- Alcantara JA, Bernarde PS, Sachett J, da Silva AM, Valente SF, Peixoto HM, et al. Stepping into a dangerous quagmire: Macroecological determinants of *Bothrops* envenoming, Brazilian Amazon. *PLoS One*. 2018; 13:e0208532. DOI:10.1371/journal.pone.0208532.
- Gutiérrez JM. Envenenamientos por mordeduras de serpiente en América Latina y el Caribe: Una visión integral de carácter regional. *Bol Malaria Salud Amb*. 2011; L1:1-16. Available from: <https://bit.ly/2R9ileR>.
- Hansson E, Sasa M, Mattison K, Robles A, Gutiérrez JM. Using geographical information systems to identify populations in need of improved accessibility to antivenom treatment for snakebite envenoming in Costa Rica. *PLoS Negl Trop Dis*. 2013; 7:e2009. DOI:10.1371/journal.pntd.0002009.
- González-Andrade F, Chippaux JP. Snake bite envenomation in Ecuador. *Trans R Soc Trop Med Hyg*. 2010; 104:588-91. DOI:10.1016/j.trstmh.2010.05.006.
- Arroyo O, Rojas G, Gutiérrez JM. Envenenamiento por mordedura de serpiente en Costa Rica en 1996: epidemiología y consideraciones clínicas. *Acta Méd Cost*. 1999; 41:23-9. Available from: <https://bit.ly/2ZzBiAN>.
- Chippaux JP, Postigo JR. Appraisal of snakebite incidence and mortality in Bolivia. *Toxicon*. 2014; 84:28-35. DOI:10.1016/j.toxicon.2014.03.007.
- De Oliveira RS, Fan HW, Sifuentes DN. Epidemiologia dos acidentes por animais peçonhentos. In: Cardoso JLC, França FOS, Wen FH, Málaque CMS, Haddad V. *Animais Peçonhentos no Brasil. Biologia, Clínica e Terapêutica dos Acidentes*. São Paulo, Brasil: Sarvier; 2009.
- Sasa M, Wasko DK, Lamar WW. Natural history of the terciopelo *Bothrops asper* (Serpentes: Viperidae) in Costa Rica. *Toxicon*. 2009; 54:904-22. DOI:10.1016/j.toxicon.2009.06.024.
- Larrick JW, Yost JA, Kaplan J. Snake bite among the Waorani Indians of Eastern Ecuador. *Trans R Soc Trop Med Hyg*. 1978; 72:542-3. DOI:10.4269/ajtmh.1991.44.93.
- Pierini SV, Warrell DA, de Paulo A, Theakston RDG. High incidence of bites and stings by snakes and other animals among rubber tappers and Amazonian Indians of the Juruá valley, Acre state, Brazil. *Toxicon*. 1996; 34:225-36. DOI:10.1016/0041-0101(95)00125-5.
- Pan American Health Organization. Policy on Ethnicity and Health. Washington, D.C.: Pan American Health Organization; 2017.
- França FOS, Málaque CMS. Accidente botrópico. In: Cardoso JLC, França FOS, Wen FH, Málaque CMS, Haddad V. *Animais Peçonhentos no Brasil. Biologia, Clínica e Terapêutica dos Acidentes*. São Paulo, Brasil: Sarvier; 2009.
- Warrell DA. Snakebites in Central and South America: Epidemiology, clinical features, and clinical management. In: Campbell J, Lamar WW. *Venomous Reptiles of the Western Hemisphere*. New York: Cornell University Press; 2004.
- Otero R, Gutiérrez J, Mesa MB, Duque E, Rodríguez O, Arango JL, et al. Complications of *Bothrops*, *Porthidium*, and *Bothriechis* snakebites in Colombia. A clinical and epidemiological study of 39 cases attended in a university hospital. *Toxicon*. 2002; 40:1107-14. DOI:10.1016/s0041-0101(02)00104-6.
- Rodríguez-Vargas A, Rodríguez-Buitrago JR, Díaz GJ. Comportamiento general de los accidentes provocados por animales venenosos en Colombia, 2006-2010. *Rev. salud pública (Bogotá)* 2012; 14:1001-09. Available from: <https://bit.ly/327m9DL>.