

Prevalence of anemia and consumption of iron-rich food groups in Mexican children and adolescents: Ensanut MC 2016

Vanessa de la Cruz-Góngora, MSc,⁽¹⁾ Salvador Villalpando, PhD,⁽²⁾ Teresa Shamah-Levy, PhD.⁽¹⁾

De la Cruz-Góngora V, Villalpando S, Shamah-Levy T. Prevalence of anemia and consumption of iron-rich food groups in Mexican children and adolescents: Ensanut MC 2016. *Salud Publica Mex* 2018;60:291-300.

<https://doi.org/10.21149/8824>

De la Cruz-Góngora V, Villalpando S, Shamah-Levy T. Prevalencia de anemia y consumo de grupos de alimentos en niños y adolescentes mexicanos: Ensanut MC 2016. *Salud Publica Mex* 2018;60:291-300.

<https://doi.org/10.21149/8824>

Abstract

Objective. To describe the prevalence of anemia and consumption of iron rich groups among Mexican children and adolescents who participated in the Halfway National Health and Nutrition Survey, 2016. **Materials and methods.** Our study sample included children and adolescents who provided full capillary hemoglobin data. Anemia was defined according to WHO criteria. Logistic regression models were used to explore the association among consumption of iron-rich food groups, sociodemographic characteristics and anemia. **Results.** In 2016, the prevalence of anemia was 26.9% in children aged 1 to 4 years old, 12.5% in those aged 5 to 11, and 9.6% in adolescents aged 12 to 19 years. Rates were the highest among females who lived in the southern and central parts of Mexico, belonged to an indigenous ethnic group and fell within the first tercile of the Household Wealth Index. Consumption of beef by preschoolers and viscera by adolescents was associated with lower risk for anemia; higher risk was associated with consumption of Liconsa milk and non-heme iron by preschoolers. **Conclusions.** Anemia is highly prevalent in Mexican children and adolescents, affecting mainly the poorest and youngest populations. Sources of heme iron are the principal dietary factor associated with low risk for anemia.

Keywords: anemia; children; adolescents; iron-rich food groups; Mexico

Resumen

Objetivo. Describir la prevalencia de anemia y el consumo de alimentos fuente de hierro en niños y adolescentes mexicanos participantes en la Encuesta Nacional de Salud y Nutrición de Medio Camino 2016. **Material y métodos.** Se analizó la información de hemoglobina capilar de niños y adolescentes. La anemia se definió con base en el criterio de la OMS. Se usaron modelos de regresión logística para identificar los grupos de alimentos y características de los individuos asociados con anemia. **Resultados.** La prevalencia de anemia fue de 26.9% en niños de 1 a 4 años; 12.5% en el grupo de 5 a 11 y 9.6% para el de 12 a 19. La mayor prevalencia de anemia se dio entre las mujeres habitantes de la región Sur y Centro, indígenas y en el primer tercil de condición de bienestar. El consumo de carne de res en preescolares y de vísceras en adolescentes se asoció con menor riesgo de anemia. El consumo de leche Liconsa y el hierro no hemo en preescolares se asoció con mayor riesgo de anemia. **Conclusiones.** La anemia es altamente prevalente en niños y adolescentes mexicanos, afectando principalmente a la población pobre y a los más jóvenes. Las fuentes dietéticas de hierro hemo permanecen como los principales factores dietéticos asociados con menor riesgo de anemia.

Palabras clave: anemia; niños; adolescentes; alimentos fuentes de hierro; México

(1) Centro de Investigación en Evaluación y Encuestas, Instituto Nacional de Salud Pública. Cuernavaca, Morelos, México.

(2) Centro de Investigación en Nutrición y Salud, Instituto Nacional de Salud Pública. Cuernavaca, Morelos, México.

Received on: May 26, 2017 • Accepted on: January 11, 2018

Corresponding author: Dr. Salvador Villalpando. Av. Universidad 655, col. Santa María Ahuacatlán. 62100, Cuernavaca, Morelos, Mexico.
E-mail:svillalp@insp.mx

Anemia persists as the main nutritional problem in the world, affecting 42.6% (95%CI 37-47) of preschool children (273 million).¹ According to the Global Burden of Disease Study 2013, the burden of anemia in developing countries accounts for 89% of all anemia-related disability.² Prevalence of anemia is trending downward, however, as a result of improved health systems, lower mortality, and broader access to health services, among other causes.² Nonetheless, despite the decline registered by Mexican surveys, prevalence remains high among the poorest and most vulnerable populations.^{3,4}

In children under 5 y,² anemia can have irreversible negative consequences for the higher intellectual functions of the brain if timely action is not taken.⁵ As for female adolescents, menstrual blood loss and increased iron requirements for growth at that age call for adequate iron-rich nutrition. Overall, anemia has been associated with increased morbidity and mortality among the most vulnerable populations.⁶

Data collected by surveys on anemia have proved insufficient to adequately analyze its principal causes and, hence, to prevent or correct its effects.³ For instance, the National Health and Nutrition Surveys (in Spanish, *Encuesta Nacional de Salud y Nutrición*, Ensanut) have clearly highlighted the role of iron deficiency (ID) as an important nutritional cause of anemia in Mexican children; however, the specific contribution of this deficiency to anemia cannot be pinpointed: among the few related biomarkers available for analysis, those that reflect ID can be underestimated due to the presence of inflammatory processes among other factors.⁷

The Mexican diet contains numerous dietary antagonists of iron absorption, such as phytates and polyphenols, owing to the high intake of corn tortillas among the population. Reports have suggested that iron intake among Mexican children and adolescents is insufficient to cover their daily requirements.⁸ Diet can thus be considered a risk factor for ID and anemia. It is noteworthy, then, that no studies have explored sources of iron by specific food groups, making it difficult to draw firm or reliable inferences for dietary policy and practice.

The aim of our study was twofold: to update existing information and provide new knowledge regarding prevalence of anemia among the Mexican children and adolescents who participated in the Halfway National Health and Nutrition Survey, 2016 (in Spanish, *Encuesta Nacional de Salud y Nutrición de Medio Camino*, Ensanut MC 2016), and to link our findings to the consumption of iron-rich food groups associated with anemia with a view to identifying opportunities for intervention.

Materials and methods

Study population. Our sample was drawn from the Ensanut MC 2016, a probabilistic, nationally representative survey with data disaggregated by regions and urban/rural strata. Analysis considered hemoglobin (Hb) data from 1 639 children aged 1 to 4.9 y; 3 181 aged 5 to 11.9 y; and 2 554 adolescents aged 12 to 19 y. Demographic and socioeconomic information was collected using ad hoc questionnaires. A detailed description of design and sampling procedures has been published elsewhere.⁹

Definition of variables

Anemia. Capillary Hb was measured using a portable HemoCue Hb photometer (HemoCue, Angelholm, Sweden). Anemia was determined when Hb concentration adjusted by altitude was <110 g/L for toddlers aged 1 to 4 y, <115g/L for children 5 to 11y, Hb<12 g/L for females aged ≥12 y and male children aged 12 to 14 y; and Hb<12g/L for males aged ≥ 15 y.¹⁰

Indigenous ethnicity was defined by whether an indigenous language was spoken at home. Localities with less than 2 500 inhabitants were considered rural, and otherwise classified as urban. A Household Wealth Index (HWI) was constructed based on household characteristics and family assets estimated through principal components analysis; the Index was divided into terciles being one as the lowest HWI score. Mexico was divided into four geographic regions: Northern, Southern, Central, and Mexico City, as in previous surveys.⁹ Anthropometric data (weight and height) were collected using validated and standardized methods.¹¹ Body mass index (BMI) and nutritional status (underweight, wasting and stunting) were computed for preschoolers according to WHO standards.^{12,13}

Children and adolescents participating in social programs were classified as Prospera and Liconsa beneficiaries.

Dietary information

Trained and standardized personnel administered a 7-day semi-quantitative food-frequency questionnaire (SFFQ) to participants, and entered the dietary data using computers in the field.¹⁴ Mothers reported the foods consumed by children in the seven days prior to the survey; adolescents responded to the questionnaire themselves in the presence of their mothers. Food consumption was calculated in grams (g) and milliliters per day. Total iron (heme and non-heme), fiber, calcium and energy were estimated using the food composi-

tion tables compiled by the National Institute of Public Health (INSP by its Spanish initials).^{*4} For the purpose of analysis, the principal iron-rich food groups were beef, pork and chicken; viscera; Liconsa milk; and beans. Corn tortillas were also taken as a relevant food group because their consumption can interfere with iron bio-availability. Respondents who reported consuming one or more food items above the 99th percentile (in grams) were excluded from analysis.

Statistical analysis

Sample characteristics and prevalence of anemia, estimated by frequency and 95% confidence intervals, were stratified by age group. Logistic regression models were constructed to test the variables associated with anemia: sex, age, BMI, HWI, nutritional status, ethnicity, rurality, geographical region, Prospera/Liconsa program affiliation, and consumption of iron-rich food groups.

All analyses were adjusted for the survey sampling design using the STATASE v14 SVY module for complex samples (College Station, USA). Statistical significance was set at $\alpha=0.05$.

The INSP Research, Ethics, and Biosecurity Committees approved the survey protocol. Prior informed consent was obtained individually from the parents of all participants.

Results

Table I illustrates the descriptive characteristics of the sample by age group.

Preschoolers

In preschoolers, the national prevalence of anemia amounted to 26.9% (95%CI 23.3, 30.9), representing an expanded population of 2.44 million anemic children. The 12 to 23-month age group yielded higher prevalence (37.9%; 95%CI 29.9, 46.7) than the 48 to 59-month age group (16%; 95%CI 11.7, 21.4), and females were affected more frequently than males ($p=0.07$). Prevalence of anemia was lower in children with overweight (OW) than in those with a normal BMI: 19.8 and 28.4%, respectively. No differences emerged between children benefiting from the Prospera/Liconsa programs and their counterparts, nor by HWI quintile, geographic

area and ethnicity (table II). Prevalence of anemia in children under five years was not statistically different from that reported by the Ensanut 2012 ($p=0.08$) (figure 1).

Consumption of iron-rich food groups

The proportion of preschool children who consumed beef and beans was higher in the non-anemic (38.2 and 82.5%, respectively) than in the anemic (26.9 and 73.2%, respectively, $p<0.05$) group (table III).

In the adjusted model, females (OR=1.5) living in the Central (OR=1.9) and Southern (OR=2.3) regions of Mexico, as well as underweight children (OR=4.0) were associated with higher odds of anemia ($p<0.05$). In contrast, older (OR=0.6) and OW (OR=0.9) children presented lower odds ($p<0.05$) for anemia (table IV). Children consuming beef (OR=0.6) and higher levels of calcium (OR=0.99) were associated with lower risk for anemia ($p<0.05$). However, those consuming non-heme iron (OR=1.8) and Liconsa milk (OR=2.1) were associated with higher risk ($p<0.05$) for anemia (table IV). Beans, tortillas and fresh fish were scarcely associated with anemia.

School-aged children

National prevalence of anemia in school-aged children was 12.5% (95%CI 10.8, 14.5), representing an expanded population of two million anemic children. Higher prevalence of anemia occurred in the five year old group (18%) compared with the 11 y group (7.5%, $p<0.001$). Children living in rural areas (15.7%) and in the South region (14.6%) yielded higher rates than their urban (11.3%) and Northern (14.6%) counterparts ($p<0.05$). Children with obesity (OB) (5%) suffered less frequently from anemia than those with a normal BMI (13.8%, $p<0.01$). Prevalence trended downward from the highest HWI quintil (p trend=0.007). No differences were observed by sex, ethnicity, or Prospera/Liconsa affiliation (table II). Prevalence in school-aged children found in our study was higher than that reported by the Ensanut 2012 ($p=0.016$) (figure 1).

Consumption of iron-rich food groups

Regarding the proportion of children who consumed iron-rich food groups, no significant differences emerged between the anemic and non-anemic groups (table III).

In the adjusted model, children who were older (OR=0.87), obese (OR=0.5), Prospera beneficiaries (OR=0.6), and in the third HWI tercile (OR=0.6) indi-

* National Institute of Public Health (INSP by its Spanish initials). Databases on the nutritional values of food compiled by INSP (unpublished), 2013.

Table I
DESCRIPTIVE CHARACTERISTICS OF MEXICAN CHILDREN AND ADOLESCENTS. MÉXICO, ENSANUT MC 2016

n (sample) N (thousands)	Preschoolers 1 639 9 093.9 Expansion			Schoolers 3 181 15 993.1 Expansion			Adolescents 2 554 18 570.7 Expansion		
	N (thousands)	%	(IC95%)	N (thousands)	%	(IC95%)	N (thousands)	%	(IC95%)
Sex (female)	4 452.5	49	(44.8-53.1)	7 901.8	49.4	(46.5-52.3)	9 117.3	49.1	(46.1-52.1)
Nutritional status									
Underweight (yes)	340.2	3.7	(2.8-4.8)	-	-	-	-	-	-
Wasting (yes)	136.0	1.5	(0.8-2.6)	-	-	-	-	-	-
Stunting (yes)	989.2	8.7	(7.6-10.1)	-	-	-	-	-	-
Body mass index (WHO)									
Normal	6 708.4	75.2	(71.4-78.7)	10 500.1	66.7	(62.5-70.5)	11 719.0	63.8	(59.9-67.5)
Overweight	2 207.2	24.8	(21.3-28.6)	2 850.7	18.1	(15.3-21.3)	4 121.5	22.4	(19.5-25.7)
Obesity	-	-	-	2 401.6	15.2	(12.3-18.7)	2 538.6	13.8	(11.3-16.8)
Area									
Urban	6 691.8	73.6	(69.7-77.1)	11 554.4	72.2	(68.1-76.0)	13 894.5	74.8	(71.1-78.2)
Rural	2 402.1	26.4	(22.9-30.3)	4 438.8	27.8	(24.0-31.9)	4 676.2	25.2	(21.8-28.9)
Geographic region									
Northern	1 811.9	19.9	(16.4-24.0)	3 436.6	21.5	(16.9-26.9)	3 476.5	18.7	(14.7-23.5)
Centre	3 216.3	35.4	(30.6-40.4)	5 226.1	32.7	(28.4-37.2)	6 516.8	35.1	(30.8-39.7)
Mexico City	1 282.5	14.1	(10.3-19.1)	2 351.6	14.7	(10.4-20.4)	2 790.0	15	(10.4-21.3)
Southern	2 783.2	30.6	(25.8-35.9)	4 978.8	31.1	(26.6-36.0)	5 787.4	31.2	(26.5-36.3)
Indigenous (yes)	128.9	1.4	(0.8-2.5)	644.9	4	(2.6-6.2)	787.0	4.2	(2.8-6.4)
Beneficiary of social program									
Prospera	1 941.2	21.3	(18.1-25.0)	4 939.6	30.9	(27.2-34.9)	6 810.7	36.7	(32.5-41.0)
Liconsa	1 068.0	11.9	(9.6-14.8)	2 111.7	13.4	(10.6-16.8)	2 114.7	11.6	(9.0-14.8)
Quintile of Household Wealth Index									
Q1	1 336.3	14.7	(12.2-17.6)	2 282.7	14.3	(11.8-17.2)	2 064.7	11.1	(8.9-13.9)
Q2	1 675.4	18.4	(15.1-22.2)	2 652.2	16.6	(14.3-19.2)	3 188.9	17.2	(14.6-20.1)
Q3	1 656.6	18.2	(15.0-22.0)	2 939.1	18.4	(16.0-21.1)	3 552.5	19.1	(16.4-22.3)
Q4	2 207.1	24.3	(20.6-28.4)	3 803.1	23.8	(19.7-28.4)	4 211.3	22.7	(19.0-26.8)
Q5	2 218.6	24.4	(19.5-30.1)	4 316.0	27	(22.3-32.2)	5 553.2	29.9	(25.7-34.5)

Ensanut MC: Encuesta Nacional de Salud y Nutrición de Medio Camino (in English, Halfway National Health and Nutrition Survey)

cated lower risk for anemia ($p<0.05$) (table I). No food-group intake was associated with anemia (table IV).

Adolescents


National prevalence of anemia in adolescents was 9.6% (95%CI 7.8, 11.8), representing an expanded population of 1.78 million anemic adolescents. Higher rates were observed among indigenous (17.6%) and female (12%) vs. non-indigenous (9.3%) and male (7.3%) adolescents

($p=0.016$). Those residing in rural (12.4%) and Southern (13.0%) localities were affected more frequently than their urban (8.7%, $p=0.06$) and Northern (6.9%, $p<0.05$) counterparts. And those with obesity (5%) registered lower prevalence than those with a normal BMI (10.5%, $p=0.006$). Prevalence trended downward in the two highest as opposed to the three lowest HWI quintiles (p trend <0.001). No differences were observed by age or Prospera/Liconsa affiliation (table II). Compared with previous surveys, current prevalence of anemia

Table II
PREVALENCE OF ANEMIA IN MEXICAN CHILDREN AND ADOLESCENTS
BY SOCIODEMOGRAPHIC CHARACTERISTICS. MÉXICO, ENSANUT MC 2016

n (sample) N (thousands)	Preschoolers 1 639 9 093.9 Expansion			Schoolers 3 181 15 993.1 Expansion			Adolescents 2 554 18 570.7 Expansion				
	N (thousands)	%	(95%CI)	N (thousands)	%	(95%CI)	N (thousands)	%	(95%CI)		
National	2 447.8	26.9	(23.3-30.9)	2 002.2	12.5	(10.8-14.5)	1 788.5	9.6	(7.8-11.8)		
Sex											
Male	1 090.0	23.5	(18.9-28.8)	1 060.3	13.1	(10.7-15.9)	691.7	7.3	(5.0-10.6)		
Female	1 357.7	30.5	(25.0-36.6)	941.9	11.9	(9.7-14.6)	1 096.9	12	(9.6-15.0)		
Age group (months)				Age group (years)			Age group (years)				
12 - 23	825.6	37.9	(29.9-46.7)	5	411.8	18.3	(13.7-24.2)	12	250.2	10.9	(7.4-15.8)
24 - 35	728.4	31.5	(24.1-40.0)	6	350.6	16.2	(12.0-21.3)	13	165.2	7.6	(4.7-12.0)
36 - 47	525.0	22.8	(17.0-29.9)	7	277.5	11.9	(7.9-17.5)	14	145.4	5.4	(3.0-9.4)
48 - 59	368.8	16	(11.7-21.4)	8	366.5	14.5	(10.0-20.5)	15	259.6	11.9	(7.7-18.0)
				9	271.4	11.5	(7.6-16.9)	16	297.7	12.6	(8.0-19.3)
				10	162.8	7.5	(4.5-12.1)	17	205.0	9.5	(6.0-14.6)
				11	161.6	7.5	(4.3-12.6)	18-19	465.4	9.9	(5.9-16.0)
Underweight											
No	2 201.3	25.4	(21.7-29.5)	-	-	-	-	-	-		
Yes	180.4	53	(35.1-70.2)	-	-	-	-	-	-		
Wasting											
No	2 289.0	26.1	(22.4-30.1)	-	-	-	-	-	-		
Yes	55.0	40.5	(16.3-70.3)	-	-	-	-	-	-		
Stunting											
No	2 049.6	25.8	(21.8-30.3)	-	-	-	-	-	-		
Yes	302.4	30.6	(22.3-40.3)	-	-	-	-	-	-		
Body mass index (WHO)											
Normal	1 906.3	28.4	(24.1-33.2)	1 451.9	13.8	(11.6-16.4)	1 228.5	10.5	(8.0-13.6)		
Overweight	437.6	19.8	(15.0-25.8)	364.1	12.8	(8.8-18.1)	402.5	9.8	(6.8-13.8)		
Obesity	-	-	-	133.4	5.6	(3.1-9.8)	125.9	5	(3.0-8.0)		
Area											
Urban	1 717.0	25.7	(21.1-30.8)	1 306.9	11.3	(9.3-13.6)	1 207.0	8.7	(6.6-11.4)		
Rural	730.8	30.4	(25.5-35.9)	695.3	15.7	(12.8-19.0)	581.6	12.4	(9.6-15.9)		
Geographic region											
Northern	323.7	17.9	(12.2-25.3)	348.5	10.1	(7.2-14.2)	238.3	6.9	(4.4-10.5)		
Centre	880.3	27.4	(21.0-34.9)	710.1	13.6	(10.7-17.1)	683.0	10.5	(7.2-15.0)		
Mexico City	302.3	23.6	(13.6-37.6)	216.0	9.2	(5.1-16.1)	113.8	4.1	(1.9-8.4)		
Southern	941.5	33.8	(28.4-39.7)	727.7	14.6	(12.0-17.7)	753.4	13	(9.7-17.3)		
Indigenous											
No	2 404.7	26.8	(23.1-30.9)	1 904.0	12.4	(10.7-14.4)	1 648.1	9.3	(7.4-11.6)		
Yes	43.0	33.4	(19.5-50.9)	98.2	15.2	(10.4-21.8)	138.4	17.6	(11.2-26.6)		
Prospera beneficiary											
No	1 846.6	25.8	(21.6-30.6)	1 401.0	12.7	(10.5-15.2)	1 081.4	9.2	(6.8-12.3)		
Yes	601.2	31	(24.5-38.2)	601.2	12.2	(9.6-15.3)	707.1	10.4	(8.0-13.4)		
Liconsa beneficiary											
No	2 228.6	28.3	(24.4-32.6)	1 697.3	12.4	(10.6-14.5)	1 628.6	10.1	(8.1-12.5)		
Yes	215.0	20.1	(13.1-29.7)	297.0	14.1	(9.4-20.6)	157.9	7.5	(4.3-12.7)		
Quintile of Household Wealth Index											
Q1	352.7	26.4	(21.0-32.6)	437.2	19.2	(14.8-24.5)	267.7	13	(9.2-18.0)		
Q2	477.5	28.5	(21.1-37.2)	338.9	12.8	(9.5-17.0)	395.6	12.4	(8.0-18.8)		
Q3	469.7	28.4	(20.9-37.2)	418.9	14.3	(10.6-18.9)	493.7	13.9	(9.0-20.8)		
Q4	614.4	27.8	(20.1-37.2)	419.8	11.0	(7.7-15.6)	292.5	6.9	(4.5-10.6)		
Q5	533.6	24.1	(16.3-34.0)	387.5	9.0	(6.0-13.2)	339.0	6.1	(4.1-9.1)		

Ensanut MC: Encuesta Nacional de Salud y Nutrición de Medio Camino (in English, Halfway National Health and Nutrition Survey)



* Statistically different from Ensanut 2012, $p < 0.05$

Ensanut: Encuesta Nacional de Salud y Nutrición (in English, National Health and Nutrition Survey)

Ensanut MC: Encuesta Nacional de Salud y Nutrición de Medio Camino (in English, Halfway National Health and Nutrition Survey)

FIGURE 1. PREVALENCE RATES OF ANEMIA IN MEXICAN CHILDREN AND ADOLESCENTS FROM 1999 TO 2016. MÉXICO, ENSANUT MC 2016

Table III
FREQUENCY OF CONSUMERS BY IRON-RICH FOOD GROUPS, AMONG NON-ANEMIC AND ANEMIC POPULATION. MÉXICO, ENSANUT MC 2016

Food group	Preschoolers		Schoolers		Adolescents	
	Non anemic % (95%CI)	Anemic % (95%CI)	Non anemic % (95%CI)	Anemic % (95%CI)	Non anemic % (95%CI)	Anemic % (95%CI)
Beef meat	38.2 (33.4-43.3)	26.9 (18.7-37.0)*	36.4 (31.8-41.4)	36.4 (29.5-43.8)	39.4 (34.9-44.0)	27.3 (19.4-37.0)*
Pork meat	30.3 (25.4-35.6)	25.1 (18.3-33.4)	38.5 (34.0-43.1)	35.8 (29.2-43.0)	38.6 (34.5-42.8)	39.8 (30.7-49.6)
Chicken	63.4 (58.3-68.2)	66.1 (58.1-73.2)	63.5 (59.1-67.7)	64.2 (57.2-70.6)	61.7 (56.6-66.5)	67.5 (57.0-75.9)
Viscera	6.2 (4.0-9.7)	7.5 (4.7-12.0)	3.0 (2.2-4.0)	3.0 (1.5-5.9)	2.00 (1.3-3.0)	0.6 (0.2-1.6)*
Fresh fish	15 (11.4-19.6)	19.1 (11.4-30.3)	15.4 (12.4-19.0)	18.5 (13.4-25.0)	14.9 (12.2-18.0)	18.8 (12.5-27.4)
Licons milk	15.9 (12.1-20.6)	14.6 (10.0-21.0)	11.4 (9.4-13.9)	12.3 (8.2-18.1)	11.7 (8.7-15.5)	8.1 (4.7-13.6)
Beans	82.8 (79.1-86.0)	73.2 (64.2-80.6)*	87.2 (84.1-89.8)	90.3 (85.3-93.7)	87.4 (85.0-89.6)	83.3 (69.9-91.4)
Tortilla	80.7 (75.4-85.1)	87 (81.8-90.8)	NA	NA	88.6 (85.5-91.1)	92.2 (87.2-95.4)

* Statistically different from non-anemic population, $p < 0.05$

Ensanut MC: Encuesta Nacional de Salud y Nutrición de Medio Camino (in English, Halfway National Health and Nutrition Survey)

in adolescents, by sex, did not differ from Ensanut 2006 results ($p=0.8$), but exceeded those of the Ensanut 2012 ($p < 0.001$) (figure 1).

Consumption of iron-rich food groups

The proportion of adolescents consuming beef and viscera was higher in the non-anemic (39.4% and 2%,

respectively) than in the anemic group (27.3 and 0.6%, respectively, $p < 0.05$) (table III).

In the adjusted model, female adolescents (OR=1.7, $p=0.029$) ranked a higher risk for anemia. Those who were obese (OR=0.5) and fell within the third HWI tercile (OR=0.5) were associated with a lower risk ($p < 0.05$). Intake of viscera (OR=0.2, $p=0.011$) was also linked to a lower risk. The association of anemia

Table IV
LOGISTIC REGRESSION MODEL FOR ANEMIA IN MEXICAN CHILDREN AND ADOLESCENTS.
MÉXICO, ENSANUT MC 2016

	Preschoolers			Schoolers			Adolescents		
	OR	(95%CI)	p	OR	(95%CI)	p	OR	(95%CI)	p
Age (years)	0.62	(0.51-0.75)	<0.001	0.87	(0.8-0.95)	0.001	1.03	(0.95-1.11)	0.487
Sex (female)	1.56	(1.08-2.26)	0.019	0.84	(0.6-1.16)	0.278	1.7	(1.06-2.75)	0.029
Rural area	1.09	(0.74-1.58)	0.671	1.27	(0.91-1.78)	0.162	1.01	(0.62-1.67)	0.955
Geographic region (northern is the reference)									
Centre	1.98	(1.06-3.7)	0.032	1.27	(0.78-2.07)	0.335	1.41	(0.74-2.71)	0.298
Mexico City and State of Mexico	1.84	(0.72-4.69)	0.201	0.81	(0.39-1.7)	0.582	0.64	(0.26-1.6)	0.34
Southern	2.27	(1.21-4.25)	0.011	1.29	(0.8-2.08)	0.288	1.53	(0.77-3.03)	0.222
Indigenous (yes)	1.53	(0.59-3.99)	0.383	1.1	(0.65-1.87)	0.728	1.85	(0.93-3.66)	0.078
Tertile of Household Wealth Index (the number 1 is the reference)									
2	1.07	(0.70-1.62)	0.757	0.9	(0.6-1.34)	0.594	1.19	(0.70-2.02)	0.51
3	1.03	(0.59-1.79)	0.925	0.57	(0.34-0.95)	0.032	0.48	(0.25-0.92)	0.027
Underweight (yes)	4.06	(1.41-11.66)	0.009	-	-	-	-	-	-
Stunting (yes)	1.07	(0.62-1.84)	0.819	-	-	-	-	-	-
Wasting (yes)	0.36	(0.06-2.30)	0.28	-	-	-	-	-	-
Body mass index (normal is the reference)									
Overweight	0.89	(0.8-0.98)	0.022	1	(0.63-1.61)	0.989	0.83	(0.54-1.29)	0.415
Obesity	-	-	-	0.46	(0.24-0.9)	0.024	0.5	(0.28-0.89)	0.019
Prospera beneficiary	1.24	(0.76-2.02)	0.379	0.65	(0.45-0.95)	0.024	0.64	(0.36-1.12)	0.117
Dietary food group consumption									
Beef	0.58	(0.39-0.88)	0.01	1.09	(0.76-1.54)	0.645	0.65	(0.39-1.09)	0.09
Pork meat	0.94	(0.57-1.55)	0.797	0.92	(0.64-1.32)	0.655	1.15	(0.75-1.75)	0.52
Chicken	1.2	(0.81-1.77)	0.365	1.28	(0.9-1.81)	0.164	1.39	(0.86-2.24)	0.183
Viscera	0.76	(0.35-1.67)	0.496	0.63	(0.25-1.59)	0.331	0.24	(0.08-0.72)	0.011
Fresh fish	1.58	(0.94-2.66)	0.083	1.31	(0.83-2.06)	0.244	1.25	(0.73-2.15)	0.409
Liconsa milk	2.1	(1.01-4.41)	0.049	1.42	(0.64-3.17)	0.391	0.57	(0.23-1.41)	0.224
Beans	0.61	(0.37-1.02)	0.061	1.23	(0.73-2.09)	0.435	0.78	(0.37-1.61)	0.492
Tortilla	1.61	(0.93-2.81)	0.09	-	NA	-	1.3	(0.63-2.70)	0.481
No heme iron (mg/d)	1.86	(1.26-2.77)	0.002	1.46	(0.89-2.42)	0.136	1.02	(0.81-1.30)	0.837
Fiber (g/d)	0.97	(0.91-1.04)	0.399	0.03	(0.01-8.72)	0.216	0.99	(0.96-1.03)	0.652
Calcium (mg/d)	0.99	(0.99-1)	<0.001	0.99	(0.99-1.00)	0.114	1.00	(1.0-1.01)	0.54
Energy (kcal/1000/day)	1.07	(0.68-1.69)	0.767	0.85	(0.68-1.07)	0.16	0.92	(0.72-1.17)	0.494
Intercept	0.25	(0.08-0.77)	0.015	0.55	(0.23-1.35)	0.191	0.03	(0.01-0.16)	<0.001

with ethnicity and beef consumption was scarcely significant (table IV).

Discussion

In Mexico, anemia affects one out of every four preschool children, one out of every seven school-aged children and one out of every 10 adolescents. Across all age groups, the prevalence trends estimated in our study matched those of the Ensanut 2006 but exceeded those of the Ensanut 2012.³

Anemia continues to afflict to the most vulnerable sectors, that is, the indigenous, poorest and youngest populations, as well as children suffering from wasting. Despite federal initiatives including Prospera¹⁵ and Liconsa¹⁶ programs—which distribute iron supplements and iron-fortified foods, respectively, to the poorest populations—anemia remains the most severe nutritional problem across all population sectors in Mexico.⁴

According to the Global Burden of Disease Study 2013, the burden of anemia showed regional differences in prevalence owing to specific and widely varying

causes. Factors as diverse as hemoglobinopathies, infectious diseases and iron-deficient diets have been found to cause anemia. In children under 5 years old, anemia is mostly due to iron-deficiency anemia (IDA).² Data from the National Health and Nutrition Examination Survey 2010 (NHANES-2010) indicated a 3.9% prevalence of anemia among children 1 to 5 years (95%CI 2.5, 5.3), ID in 7.1% (5.5, 8.7) and 1.1% (0.6, 1.7) in IDA.¹⁷ In Mexican children, previous surveys have highlighted nutritional deficiencies as one of its major causes.¹⁸ Our study was limited by the lack of biomarker measurements for identifying total body iron, Vitamin-B12, folate, CRP and hepcidin. Such data would have allowed us to pinpoint the proportion of deficiencies in iron, vitamins and other minerals, as well as the presence of inflammation, as causes of anemia. In turn, this would have served to determine their contributions to the high prevalence rates reported by the Ensanut MC 2016.

Contrary to expectations, our findings indicated lower risk for anemia in preschoolers with OW, as well as in children and adolescents with obesity (OB). It has been documented that OW/OB are emerging risk factors for ID given their pro-inflammatory role in promoting increased hepcidin synthesis and restricted iron export activity in cells.^{7,19,20} While ID resulting from an inflammatory process (commonly promoted by OB) can lead to anemia of inflammation, cross-sectional studies of Mexican children have consistently demonstrated that a high BMI is inversely associated with anemia.^{3,21} García and colleagues found no association between low-iron status and OW/OB in a sample of Mexican children.²² In contrast, Cepeda-López and colleagues documented that OB was a predictor for low-iron status among Mexican children.²⁰ Divergent results can be explained by the use of different biomarkers in defining ID. In Mexico, it is noteworthy that no studies have analyzed the association between BMI and anemia. Because cross-sectional designs allow for limited understanding as to how BMI trajectories influence Hb concentration over time, it is recommended that future studies explore such an association using a cohort design. Given the double burden of nutrition and the epidemiological transition Mexico has been facing, the dietary patterns of subjects with OW may be marked by greater vitamin and mineral density coupled with higher energy intake. In turn, this could be leading to higher Hb concentrations and lower prevalence of anemia in comparison with normal BMI subjects. Additional factors including a higher socioeconomic status (SES) in subjects with OW may also explain such an association. We did not explore SES data by statistical interaction; further analysis should be undertaken,

stratifying by BMI, in order to better understand the ways in which SES, dietary iron absorption, adiposity and inflammation influence anemia among Mexican children and adolescents.

While dietary analysis under the Ensanut 2012 showed that iron inadequacy was highly prevalent among Mexican children and adolescents,⁸ it did not clarify whether this condition was associated with anemia. Given the dearth of studies using food groups to investigate such an association in Mexico, we decided to analyze the food groups richest in iron in an effort to elucidate the relationship between iron inadequacy and anemia. Our results indicated that beef and viscera, well-known sources of bioavailable heme iron, were associated with lower risk for anemia. Preschoolers exhibited the lowest risk, followed by adolescents. The highest association, found in adolescent consumers of viscera, indicated 77% lower odds of anemia compared to non-consumers. Intake of these foods has been widely recognized to improve Hb concentrations; however, only a small proportion of Mexican children and adolescents consume beef. This suggests that certain population groups (from non-indigenous, urban and higher SES sectors - data not shown) enjoy access to animal foods while others (from indigenous and poverty-ridden contexts) do not, with the latter living in conditions of high food insecurity.²³ Chicken was the most frequently consumed meat across all age groups. Although proven to be an abundant source of iron,²⁴ it was not associated with anemia. More comprehensive studies are required in order to gain a better understanding of results obtained thus far. Analysis should consider the amounts, types and frequency of meat consumed, use a 24-hour recall method, and establish a clear contrast between anemic and non-anemic populations. Our study aimed at identifying the association of iron-rich food groups with anemia.

According to our findings, food groups rich in non-heme iron such as beans and legumes were consumed extensively throughout the population, with intake declining, however, among anemic preschoolers: Prevalence of anemia among preschool consumers of beans was eight percentage points higher (33%) than among non-consumers in the same age group (25%, $p=0.027$) (data not shown). Intake was not different in other age groups; however, school-aged children yielded the highest proportion of bean consumers. This suggests that beans combined with other foods that facilitate iron absorption may be an effective dietary source of iron for preventing anemia.

Although other non-heme iron food groups were not evaluated in our study, we were able to identify that

total dietary non-heme iron was associated with higher risk for anemia among preschoolers.

The iron-rich food groups associated with anemia in this study are also sources of other micronutrients that contribute to Hb synthesis such as Vitamin A, Vitamin B12, and zinc. The food-synergy approach may therefore be useful for understanding the interaction of diet with risk for anemia.^{25,26}

The present study is subject to several limitations. First, a cross-sectional design rendered reverse causality an alternative explanation as regards the association of food groups with anemia, despite adjustment for confounders. Second, misclassification of food groups, under- and over-reporting of food consumption (specifically by subjects with OB and low body weight), and use of a food frequency questionnaire may have undermined accuracy in estimating the association between food group consumption and anemia. Third, changes in population-level determinants of anemia, particularly dietary intake, can influence trends. The upswing in prevalence over the past three years was unexpected. Although the three Ensanut consulted used the same method for measuring Hb and classifying anemia, and population characteristics remained unchanged during the period analyzed, we were unable to identify the underlying factors of increased prevalence. Methodological issues concerning Ensanut MC 2016 include seasonality for data collection.

The major *strength* of our *study* resides in its nationally representative samples of children and adolescents. Additionally, in a random subsample of this population, capillary Hb and venous Hb measured by means of a hematology analyzer yielded results that were consistently reliable, ruling out potential measurement bias. Our results provide updated evidence of the prevalence of anemia for policy makers, thus improving opportunities for interventions.

All of these findings pinpoint anemia as a serious challenge for public health nutrition, despite a number of efforts by the government to prevent it, especially in the vulnerable populations. The current nutrition policy relying on social programs such as Prospera and Liconsa should reinforce micronutrient supplementation and education for mothers of childbearing age. Studies that address the main causes of anemia more comprehensively in Mexican children are required as a basis for more targeted public health policies.

In conclusion, anemia is highly prevalent among Mexican children and adolescents, affecting mainly indigenous peoples as well as the youngest and poorest populations. Dietary sources of bioavailable iron remain as the main dietary factor associated with lower risk for anemia. Further analyses taking the whole

diet as well as other demographic characteristics into account are necessary for improving dietary recommendations designed specifically for the vulnerable populations.

Declaration of conflict of interests. The authors declare that they have no conflict of interests.

References

1. World Health Organization. The Global Prevalence of Anemia in 2011. Geneva: WHO, 2015 [cited may, 2017]. Available from: http://apps.who.int/iris/bitstream/10665/1177094/1/9789241564960_eng.pdf?ua=1
2. Kassebaum NJ. The Global Burden of Anemia. *Hematol Oncol Clin North Am.* 2016;30(2):247-308. <https://doi.org/10.1016/j.hoc.2015.11.002>
3. De La Cruz-Góngora V, Villalpando S, Mundo-Rosas V, Shamah-Levy T. Prevalencia de anemia en niños y adolescentes mexicanos: Comparativo de tres encuestas nacionales. *Salud Publica Mex.* 2013;55(Suppl 2):180-9.
4. Gutiérrez JP, Rivera-Dommarco JA, Shamah-Levy T, Villalpando-Hernández S, Franco A, Cuevas-Nasu L, et al. Encuesta Nacional de Salud y Nutrición 2012. Resultados Nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2012 [cited may, 2017]. Available from: <http://ensanut.insp.mx/informes/ENSANUT2012ResultadosNacionales.pdf>
5. Hermoso M, Vucic V, Vollhardt C, Arsic A, Roman-Viñas B, Iglesia-Altaba I, et al. The effect of iron on cognitive development and function in infants, children and adolescents: A systematic review. *Ann Nutr Metab.* 2011;59(2-4):154-65. <https://doi.org/10.1159/000334490>
6. Tolentino K, Friedman JF. An update on anemia in less developed countries. *Am J Trop Med Hyg.* 2007;77(1):44-51.
7. Nemeth E, Valore EV, Territo M, Schiller G, Lichtenstein A, Ganz T. Hepcidin, a putative mediator of anemia of inflammation, is a type II acute-phase protein. *Blood.* 2003;101(7):2461-3. <https://doi.org/10.1182/blood-2002-10-3235>
8. Sánchez-Pimienta TG, López-Olmedo N, Rodríguez-Ramírez S, García-Guerra A, Rivera JA, Carriquiry AL, et al. High Prevalence of Inadequate Calcium and Iron Intakes by Mexican Population Groups as Assessed by 24-Hour Recalls. *J Nutr.* 2016;146(9):1874S-80S. <https://doi.org/10.3945/jn.115.227074>
9. Romero-Martínez M, Shamah-Levy T, Cuevas-Nasu L, Méndez Gómez-Humarán I, Gaona-Pineda EB, Gómez-Acosta LM, et al. Diseño metodológico de la Encuesta Nacional de Salud y Nutrición de Medio Camino 2016. *Salud Publica Mex.* 2017;59(3):299-305. <https://doi.org/10.21149/8593>
10. World Health Organization. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. Vitamin and Mineral Nutrition Information System. Geneva, Switzerland: WHO, 2011 [cited January, 2015]. Available from: <http://www.who.int/vmnis/indicators/haemoglobin.pdf>
11. Lohman TG, Roche AF, Martorell R. Anthropometric standardization reference manual: Abridged edition. Champaign, Ill: Hum Kinet Books, 1988;177.
12. World Health Organization. Interpretation Guide Nutrition Landscape Information System (NLIS). Geneva: WHO, 2010.
13. World Health Organization. WHO child growth standards: length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: methods and development. Geneva: WHO, 2006 [cited April, 2016]. Available from: http://www.who.int/childgrowth/standards/Technical_report.pdf
14. Ramírez-Silva I, Jiménez-Aguilar A, Valenzuela-Bravo D, Martínez-Tapia B, Rodríguez-Ramírez S, Gaona-Pineda EB, et al. Methodology for analysis of dietary information from Semi-quantitative Food Frequency Questionnaire of the Mexican National Health and Nutrition Survey 2012. *Salud Publica Mex.* 2016;58(6):629-38. <https://doi.org/10.21149/spm.v58i6.7974>

15. Rivera JA, Sotres-Alvarez D, Habicht JP, Shamah T, Villalpando S. Impact of the Mexican Program for Education, Health, and Nutrition (Progresa) on Rates of Growth and Anemia in Infants and Young Children. *JAMA*. 2004;291(21):2563. <https://doi.org/10.1001/jama.291.21.2563>
16. Rivera JA, Shamah T, Villalpando S, Monterrubio E. Effectiveness of a large-scale iron-fortified milk distribution program on anemia and iron deficiency in low-income young children in Mexico. *Am J Clin Nutr*. 2010;91(2):431-9. <https://doi.org/10.3945/ajcn.2009.28104>
17. Gupta P, Perrine C, Mei Z, Scanlon K. Iron, Anemia, and Iron Deficiency Anemia among Young Children in the United States. *Nutrients*. 2017;9(8):876.
18. De la Cruz-Góngora V, Villalpando S, Rebollar R, Shamah-Levy T, Méndez-Gómez Humarán I. Nutritional causes of anemia in Mexican children under 5 years. Results from the 2006 National Health and Nutrition Survey. *Salud Publica Mex*. 2012;54(2):108-15. <https://doi.org/10.1590/S0036-36342012000200006>
19. Hutchinson C. A review of iron studies in overweight and obese children and adolescents: a double burden in the young? *Eur J Nutr*. 2016;55(7):2179-97. <https://doi.org/10.1007/s00394-016-1155-7>
20. Cepeda-Lopez AC, Osendarp SJM, Melse-Boonstra A, Aeberli I, Gonzalez-Salazar F, Feskens E, et al. Sharply higher rates of iron deficiency in obese Mexican women and children are predicted by obesity-related inflammation rather than by differences in dietary iron intake. *Am J Clin Nutr*. 2011;93(5):975-83. <https://doi.org/10.3945/ajcn.110.005439>
21. Syed S, Addo OY, De La Cruz-Góngora V, Ashour FAS, Ziegler TR, Suchdev PS. Determinants of anemia among school-aged children in Mexico, the United States and Colombia. *Nutrients*. 2016;8(7):387. <https://doi.org/10.3390/nu8070387>
22. García OP, Ronquillo D, Caamaño MC, Martínez G, Camacho M, López V, et al. Zinc, iron and vitamins A, C and E are associated with obesity, inflammation, lipid profile and insulin resistance in Mexican school-aged children. *Nutrients*. 2013;5(12):5012-30. <https://doi.org/10.3390/nu5125012>
23. Shamah-Levy T, Villalpando-Hernández S, Cuevas-Nassu L, Gaona-Pineda EB, Gómez-Humarán I, Rivera-Dommarco JA. Relationship of household food insecurity and anemia in children receiving the benefit from food assistance programs in San Luis Potosí, Mexico. *FASEB J*. 2015;29(Suppl 1):S585-S588.
24. Marangoni F, Corsello G, Cricelli C, Ferrara N, Ghiselli A, Lucchin L, et al. Role of poultry meat in a balanced diet aimed at maintaining health and wellbeing: An Italian consensus document. *Food Nutr Res*. 2015;59:1-11. <https://doi.org/10.3402/fnr.v59.27606>
25. Tapsell LC, Neale EP, Satija A, Hu FB. Foods, Nutrients, and Dietary Patterns: Interconnections and Implications for Dietary Guidelines. *Adv Nutr An Int Rev J*. 2016;7(3):445-54. <https://doi.org/10.3945/an.115.011718>
26. Jacobs DR, Steffen LM. Nutrients, foods, and dietary patterns as exposures in research: A framework for food synergy. *Am J Clin Nutr*. 2003;78(Suppl 3):508S-513S. <https://doi.org/10.1093/ajcn/78.3.508S>